

PTGS2 Antibody (Center P378)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11553C

Specification

PTGS2 Antibody (Center P378) - Product Information

Application	WB, IHC-P, IF,E
Primary Accession	P35354
Other Accession	NP_000954.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	363-391

PTGS2 Antibody (Center P378) - Additional Information

Gene ID 5743

Other Names

Prostaglandin G/H synthase 2, Cyclooxygenase-2, COX-2, PHS II, Prostaglandin H2 synthase 2, PGH synthase 2, PGHS-2, Prostaglandin-endoperoxide synthase 2, PTGS2, COX2

Target/Specificity

This PTGS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 363-391 amino acids from the Central region of human PTGS2.

Dilution

WB~~1:2000
IHC-P~~1:10~50
IF~~1:10~50
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PTGS2 Antibody (Center P378) is for research use only and not for use in diagnostic or therapeutic procedures.

PTGS2 Antibody (Center P378) - Protein Information

Name PTGS2 ([HGNC:9605](#))

Function Dual cyclooxygenase and peroxidase in the biosynthesis pathway of prostanoids, a class of C20 oxylipins mainly derived from arachidonate ((5Z,8Z,11Z,14Z)-eicosatetraenoate, AA, C20:4(n-6)), with a particular role in the inflammatory response (PubMed:[11939906](#), PubMed:[16373578](#), PubMed:[19540099](#), PubMed:[22942274](#), PubMed:[26859324](#), PubMed:[27226593](#), PubMed:[7592599](#), PubMed:[7947975](#), PubMed:[9261177](#)). The cyclooxygenase activity oxygenates AA to the hydroperoxy endoperoxide prostaglandin G2 (PGG2), and the peroxidase activity reduces PGG2 to the hydroxy endoperoxide prostaglandin H2 (PGH2), the precursor of all 2-series prostaglandins and thromboxanes (PubMed:[16373578](#), PubMed:[22942274](#), PubMed:[26859324](#), PubMed:[27226593](#), PubMed:[7592599](#), PubMed:[7947975](#), PubMed:[9261177](#)). This complex transformation is initiated by abstraction of hydrogen at carbon 13 (with S-stereochemistry), followed by insertion of molecular O₂ to form the endoperoxide bridge between carbon 9 and 11 that defines prostaglandins. The insertion of a second molecule of O₂ (bis-oxygenase activity) yields a hydroperoxy group in PGG2 that is then reduced to PGH2 by two electrons (PubMed:[16373578](#), PubMed:[22942274](#), PubMed:[26859324](#), PubMed:[27226593](#), PubMed:[7592599](#), PubMed:[7947975](#), PubMed:[9261177](#)). Similarly catalyzes successive cyclooxygenation and peroxidation of dihomo-gamma-linoleate (DGLA, C20:3(n-6)) and eicosapentaenoate (EPA, C20:5(n-3)) to corresponding PGH1 and PGH3, the precursors of 1- and 3-series prostaglandins (PubMed:[11939906](#), PubMed:[19540099](#)). In an alternative pathway of prostanoid biosynthesis, converts 2-arachidonoyl lysophospholipids to prostanoid lysophospholipids, which are then hydrolyzed by intracellular phospholipases to release free prostanoids (PubMed:[27642067](#)). Metabolizes 2-arachidonoyl glycerol yielding the glyceryl ester of PGH2, a process that can contribute to pain response (PubMed:[22942274](#)). Generates lipid mediators from n-3 and n-6 polyunsaturated fatty acids (PUFAs) via a lipoxygenase-type mechanism. Oxygenates PUFAs to hydroperoxy compounds and then reduces them to corresponding alcohols (PubMed:[11034610](#), PubMed:[11192938](#), PubMed:[9048568](#), PubMed:[9261177](#)). Plays a role in the generation of resolution phase interaction products (resolvins) during both sterile and infectious inflammation (PubMed:[12391014](#)). Metabolizes docosahexaenoate (DHA, C22:6(n-3)) to 17R-HDHA, a precursor of the D-series resolvins (RvDs) (PubMed:[12391014](#)). As a component of the biosynthetic pathway of E-series resolvins (RvEs), converts eicosapentaenoate (EPA, C20:5(n-3)) primarily to 18S-HEPE that is further metabolized by ALOX5 and LTA4H to generate 18S-RvE1 and 18S-RvE2 (PubMed:[21206090](#)). In vascular endothelial cells, converts docosapentaenoate (DPA, C22:5(n-3)) to 13R-HDPA, a precursor for 13-series resolvins (RvTs) shown to activate macrophage phagocytosis during bacterial infection (PubMed:[26236990](#)). In activated leukocytes, contributes to oxygenation of hydroxyeicosatetraenoates (HETE) to diHETES (5,15-diHETE and 5,11-diHETE) (PubMed:[22068350](#), PubMed:[26282205](#)). Can also use linoleate (LA, (9Z,12Z)-octadecadienoate, C18:2(n-6)) as substrate and produce hydroxyoctadecadienoates (HODEs) in a regio- and stereospecific manner, being (9R)-HODE ((9R)-hydroxy-(10E,12Z)-octadecadienoate) and (13S)-HODE ((13S)-hydroxy-(9Z,11E)-octadecadienoate) its major products (By similarity). During neuroinflammation, plays a role in neuronal secretion of specialized preresolving mediators (SPMs) 15R-lipoxin A4 that regulates phagocytic microglia (By similarity).

Cellular Location

Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein. Nucleus inner membrane; Peripheral membrane protein. Nucleus outer membrane; Peripheral membrane protein. Note=Detected on the luminal side of the endoplasmic reticulum and nuclear envelope


PTGS2 Antibody (Center P378) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PTGS2 Antibody (Center P378) - Images




Confocal immunofluorescent analysis of PTGS2 Antibody (Center P378) (Cat. #AP11553c) with A549 cell followed by Alexa Fluor® 489-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



PTGS2 Antibody (Center P378) (Cat. #AP11553c) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the PTGS2 antibody detected the PTGS2 protein (arrow).



Anti-PTGS2 Antibody (Center P378) at 1:2000 dilution + HACAT whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 67 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



PTGS2 Antibody (Center P378) (Cat. #AP11553c) immunohistochemistry analysis in formalin fixed and paraffin embedded human uterus tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of PTGS2 Antibody (Center P378) for immunohistochemistry. Clinical relevance has not been evaluated.

PTGS2 Antibody (Center P378) - Background

Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isoforms of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isoform. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis.

PTGS2 Antibody (Center P378) - References

Duggan, K.C., et al. J. Biol. Chem. 285(45):34950-34959(2010)
Feher, A., et al. Am J Geriatr Psychiatry 18(11):983-987(2010)
Wang, C.H., et al. Anticancer Res. 30(9):3649-3653(2010)

Han, E.H., et al. J. Toxicol. Environ. Health Part A 73 (21-22), 1451-1464 (2010) :

Cao, H., et al. Tohoku J. Exp. Med. 222(1):15-21(2010)

PTGS2 Antibody (Center P378) - Citations

- [Predictive value of APAF-1 and COX-2 expression in pathologic complete response to neoadjuvant chemoradiotherapy for patients with locally advanced rectal adenocarcinoma.](#)