

OR52I2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP11831b**Specification**

OR52I2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q8NH67
Other Accession	NP_001005170.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	38353
Antigen Region	305-333

OR52I2 Antibody (C-term) - Additional Information**Gene ID** 143502**Other Names**

Olfactory receptor 52I2, Olfactory receptor OR11-12, OR52I2

Target/Specificity

This OR52I2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 305-333 amino acids from the C-terminal region of human OR52I2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR52I2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR52I2 Antibody (C-term) - Protein Information**Name** OR52I2**Function** Odorant receptor.


Cellular Location

Cell membrane; Multi-pass membrane protein.

OR52I2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR52I2 Antibody (C-term) - Images

OR52I2 Antibody (C-term) (Cat. #AP11831b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the OR52I2 antibody detected the OR52I2 protein (arrow).

OR52I2 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR52I2 Antibody (C-term) - References

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)
Vanti, W.B., et al. Biochem. Biophys. Res. Commun. 305(1):67-71(2003)