

**UGT2B15 Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP12375C**

**Specification**

---

**UGT2B15 Antibody (Center) - Product Information**

Application	IHC-P, WB, FC, IF,E
Primary Accession	<a href="#">P54855</a>
Other Accession	<a href="#">NP_001067.2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	156-185

**UGT2B15 Antibody (Center) - Additional Information**

**Gene ID** 7366

**Other Names**

UDP-glucuronosyltransferase 2B15, UDPGT 2B15, HLUG4, UDP-glucuronosyltransferase 2B8, UDPGT 2B8, UDPGTh-3, UGT2B15, UGT2B8

**Target/Specificity**

This UGT2B15 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 156-185 amino acids from the Central region of human UGT2B15.

**Dilution**

IHC-P~~1:100  
WB~~1:500  
FC~~1:10~50  
IF~~1:10~50  
E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

UGT2B15 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**UGT2B15 Antibody (Center) - Protein Information**

**Name** UGT2B15 ([HGNC:12546](#))

**Function** UDP-glucuronosyltransferase (UGT) that catalyzes phase II biotransformation reactions in which lipophilic substrates are conjugated with glucuronic acid to increase the metabolite's water solubility, thereby facilitating excretion into either the urine or bile (PubMed:[16595710](#), PubMed:[18719240](#), PubMed:[23288867](#), PubMed:[7835232](#), PubMed:[9295060](#)). Essential for the elimination and detoxification of drugs, xenobiotics and endogenous compounds (PubMed:[7835232](#)). Catalyzes the glucuronidation of endogenous steroid hormones such as androgens (testosterone, androsterone) and estrogens (estradiol, epiestradiol, estriol, catechol estrogens) (PubMed:[16595710](#), PubMed:[18719240](#), PubMed:[23288867](#), PubMed:[7835232](#), PubMed:[9295060](#)). Displays glucuronidation activity toward several classes of xenobiotic substrates, including phenolic compounds (eugenol, 4-nitrophenol, 4-hydroxybiphenyl) and phenylpropanoids (naringenin, coumarins) (PubMed:[7835232](#)). Catalyzes the glucuronidation of monoterpene alcohols such as borneol, menthol and isomenthol, a class of natural compounds used in essential oils (By similarity).

**Cellular Location**

Endoplasmic reticulum membrane; Single-pass membrane protein


**Tissue Location**

Expressed in many tissues. Present in liver, prostate and testis.


**UGT2B15 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**UGT2B15 Antibody (Center) - Images**


Confocal immunofluorescent analysis of UGT2B15 Antibody (Center) (Cat#AP12375c) with NCI-H460 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Anti-UGT2B15 Antibody (Center) at 1:500 dilution + Human liver tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 61 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of AP12375C on paraffin-embedded Human liver tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



UGT2B15 Antibody (Center) (Cat. #AP12375c) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **UGT2B15 Antibody (Center) - Background**

The UGTs are of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds. UGT2B8 demonstrates reactivity with estriol. See UGT2B4 (MIM 600067).

#### **UGT2B15 Antibody (Center) - References**

Yong, M., et al. Cancer Epidemiol. Biomarkers Prev. 19(2):537-546(2010)  
Sun, C., et al. Hum. Mutat. 31(1):99-107(2010)  
Ross, C.J., et al. Nat. Genet. 41(12):1345-1349(2009)  
He, X., et al. Br J Clin Pharmacol 68(5):721-730(2009)  
Holmes, M.V., et al. PLoS ONE 4 (12), E7960 (2009) :

#### **UGT2B15 Antibody (Center) - Citations**

- [Steroidogenic enzyme expression in the human fetal liver and potential role in the endocrinology of pregnancy.](#)