

Sestrin-3 Antibody (Center Y335)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP12471c**Specification**

Sestrin-3 Antibody (Center Y335) - Product Information

Application	IHC-P, WB,E
Primary Accession	P58005
Other Accession	Q9CYP7 , NP_653266.2
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	320-347

Sestrin-3 Antibody (Center Y335) - Additional Information**Gene ID** 143686**Other Names**

Sestrin-3, SESN3, SEST3

Target/Specificity

This Sestrin-3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 320-347 amino acids from the Central region of human Sestrin-3.

Dilution

IHC-P~~1:10~50

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Sestrin-3 Antibody (Center Y335) is for research use only and not for use in diagnostic or therapeutic procedures.

Sestrin-3 Antibody (Center Y335) - Protein Information**Name** SESN3 ([HGNC:23060](#))

Function May function as an intracellular leucine sensor that negatively regulates the TORC1 signaling pathway (PubMed:[25263562](#)). May also regulate the insulin-receptor signaling pathway through activation of TORC2 (By similarity). This metabolic regulator may also play a role in protection against oxidative and genotoxic stresses (By similarity). May prevent the accumulation of reactive oxygen species (ROS) through the alkylhydroperoxide reductase activity born by the N-terminal domain of the protein (By similarity).

Cellular Location

Cytoplasm.

Tissue Location


Widely expressed..

Sestrin-3 Antibody (Center Y335) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Sestrin-3 Antibody (Center Y335) - Images



Sestrin-3 Antibody (Center Y335) (Cat. #AP12471c) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the Sestrin-3 antibody detected the Sestrin-3 protein (arrow).



Anti-Sestrin-3 Antibody (Center Y335) at 1:2000 dilution + LNCaP whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Sestrin-3 Antibody (Center Y335) (Cat. #AP12471c) immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of Sestrin-3 Antibody (Center Y335) for immunohistochemistry. Clinical relevance has not been evaluated.

Sestrin-3 Antibody (Center Y335) - Background

Sestrins form a small family of redox enzymes which regulate accumulation of Reactive Oxygen Species. They protect cells and their DNA against oxidative damage and regulate cell growth and viability. These genes are often deregulated in cancers and their inactivation accelerates the growth of model tumors in mice.

Sestrin-3 Antibody (Center Y335) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Chen, C.C., et al. Dev. Cell 18(4):592-604(2010)
Peeters, H., et al. Hum. Genet. 112 (5-6), 573-580 (2003) :

Sestrin-3 Antibody (Center Y335) - Citations

- [ATF4-Mediated Upregulation of REDD1 and Sestrin2 Suppresses mTORC1 Activity during Prolonged Leucine Deprivation](#)
- [Evidence for a Role for Sestrin1 in Mediating Leucine-Induced Activation of mTORC1 in Skeletal Muscle](#)
- [Leucine induced dephosphorylation of Sestrin2 promotes mTORC1 activation](#)
- [GCN2 sustains mTORC1 suppression upon amino acid deprivation by inducing Sestrin2](#)
- [Amino Acid-Induced Activation of mTORC1 in Rat Liver Is Attenuated by Short-Term Consumption of a High-Fat Diet](#)