

# DNAJB11 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12621a

## Specification

# **DNAJB11** Antibody (N-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region WB, IHC-P,E <u>O9UBS4</u> <u>O6TUG0</u>, <u>O99KV1</u>, <u>O3ZBA6</u>, <u>NP\_057390.1</u> Human Bovine, Mouse, Rat Rabbit Polyclonal Rabbit IgG 40514 63-92

## **DNAJB11** Antibody (N-term) - Additional Information

Gene ID 51726

### **Other Names**

DnaJ homolog subfamily B member 11, APOBEC1-binding protein 2, ABBP-2, DnaJ protein homolog 9, ER-associated DNAJ, ER-associated Hsp40 co-chaperone, Endoplasmic reticulum DNA J domain-containing protein 3, ER-resident protein ERdj3, ERdj3, ERj3p, HEDJ, Human DnaJ protein 9, hDj-9, PWP1-interacting protein 4, DNAJB11, EDJ, ERJ3, HDJ9

## Target/Specificity

This DNAJB11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 63-92 amino acids from the N-terminal region of human DNAJB11.

**Dilution** WB~~1:1000 IHC-P~~1:10~50 E~~Use at an assay dependent concentration.

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## Precautions

DNAJB11 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## **DNAJB11** Antibody (N-term) - Protein Information



Name DNAJB11

Synonyms EDJ, ERJ3, HDJ9

**Function** As a co-chaperone for HSPA5 it is required for proper folding, trafficking or degradation of proteins (PubMed:<u>10827079</u>, PubMed:<u>15525676</u>, PubMed:<u>29706351</u>). Binds directly to both unfolded proteins that are substrates for ERAD and nascent unfolded peptide chains, but dissociates from the HSPA5-unfolded protein complex before folding is completed (PubMed:<u>15525676</u>). May help recruiting HSPA5 and other chaperones to the substrate. Stimulates HSPA5 ATPase activity (PubMed:<u>10827079</u>). It is necessary for maturation and correct trafficking of PKD1 (PubMed:<u>29706351</u>).

**Cellular Location** Endoplasmic reticulum lumen Note=Associated with the ER membrane in a C-terminally epitope-tagged construct


**Tissue Location** Widely expressed.

## **DNAJB11** Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>


## DNAJB11 Antibody (N-term) - Images



Western blot analysis of lysate from SW480 cell line, using DNAJB11 Antibody (N-term)(Cat. #AP12621a).AP12621a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at



1:5000 dilution was used as the secondary antibody.Lysate at 35ug per lane.



DNAJB11 Antibody (N-term) (Cat. #AP12621a)immunohistochemistry analysis in formalin fixed and paraffin embedded human placenta tissue followed by peroxidase conjugation of the secondary antibody and DAB staining.This data demonstrates the use of DNAJB11 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

# DNAJB11 Antibody (N-term) - Background

DNAJB11 belongs to the evolutionarily conserved DNAJ/HSP40 family of proteins, which regulate molecular chaperone activity by stimulating ATPase activity. DNAJ proteins may have up to 3 distinct domains: a conserved 70-amino acid J domain, usually at the N terminus; a glycine/phenylalanine (G/F)-rich region; and a C-terminal cysteine-rich region (Ohtsuka and Hata, 2000 [PubMed 11147971]).

# DNAJB11 Antibody (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Wen, K.W., et al. Oncogene 29(24):3532-3544(2010) Bernal-Bayard, J., et al. J. Biol. Chem. 285(21):16360-16368(2010) Vembar, S.S., et al. J. Biol. Chem. 284(47):32462-32471(2009) Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)