

**CYP2S1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP12635B**

### Specification

#### CYP2S1 Antibody (C-term) - Product Information

Application	WB, IF, IHC-P,E
Primary Accession	<a href="#">Q96SQ9</a>
Other Accession	<a href="#">NP_085125.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	399-428

#### CYP2S1 Antibody (C-term) - Additional Information

##### Gene ID 29785

##### Other Names

Cytochrome P450 2S1, CYPIIS1, CYP2S1

##### Target/Specificity

This CYP2S1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 399-428 amino acids from the C-terminal region of human CYP2S1.

##### Dilution

WB~~1:1000

IF~~1:10~50

IHC-P~~1:10~50

E~~Use at an assay dependent concentration.

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

CYP2S1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### CYP2S1 Antibody (C-term) - Protein Information

Name CYP2S1 {ECO:0000303|PubMed:11181079, ECO:0000312|HGNC:HGNC:15654}

**Function** A cytochrome P450 monooxygenase involved in the metabolism of retinoids and eicosanoids (PubMed:[12711469](#), PubMed:[21068195](#)). In epidermis, may contribute to the oxidative metabolism of all-trans- retinoic acid. For this activity, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:[12711469](#)). Additionally, displays peroxidase and isomerase activities toward various oxygenated eicosanoids such as prostaglandin H2 (PGH2) and hydroperoxyeicosatetraenoates (HPETEs) (PubMed:[21068195](#)). Independently of cytochrome P450 reductase, NADPH, and O2, catalyzes the breakdown of PGH2 to hydroxyheptadecatrienoic acid (HHT) and malondialdehyde (MDA), which is known to act as a mediator of DNA damage (PubMed:[21068195](#)).

#### Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

#### Tissue Location

Expressed at higher levels in extrahepatic tissues including trachea, lung, stomach, small intestine, colon, kidney, breast, placenta and spleen (PubMed:11181079, PubMed:12711469)


Expressed in peripheral blood leukocytes (PubMed:11181079) Constitutively expressed in skin (at protein level) (PubMed:12711469)

### CYP2S1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CYP2S1 Antibody (C-term) - Images




Confocal immunofluorescent analysis of CYP2S1 Antibody (C-term)(Cat#AP12635b) with HepG2


cell followed by Alexa Fluor®488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



CYP2S1 Antibody (C-term) (Cat. #AP12635b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the CYP2S1 antibody detected the CYP2S1 protein (arrow).



Western blot analysis of lysate from BEAS 2B cell line, using CYP2S1 Antibody (C-term)(Cat. #AP12635b). AP12635b was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



CYP2S1 Antibody (C-term) (Cat. #AP12635b) immunohistochemistry analysis in formalin fixed and paraffin embedded human stomach tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of CYP2S1 Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.

#### **CYP2S1 Antibody (C-term) - Background**

This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum. In rodents, the homologous protein has been shown to metabolize certain carcinogens; however, the specific function of the human protein has not been determined.

#### **CYP2S1 Antibody (C-term) - References**

Bui, P.H., et al. Mol. Pharmacol. 76(5):1031-1043(2009) Saito, A., et al. J. Hum. Genet. 54(6):317-323(2009) Deb, S., et al. Expert Opin Drug Metab Toxicol 5(4):367-380(2009) Jang, Y.J., et al. Ther Drug Monit 29(3):292-298(2007) Hanzawa, Y., et al. Drug Metab. Pharmacokinet. 22(2):136-140(2007)

#### **CYP2S1 Antibody (C-term) - Citations**

- [Upregulation of CYP2S1 by oxaliplatin is associated with p53 status in colorectal cancer cell lines.](#)
- [CYP2S1 Depletion Enhances Colorectal Cell Proliferation is Associated with PGE2-Mediated Activation of  \$\beta\$ -catenin Signaling.](#)