

RPL15 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP12772A

Specification

RPL15 Antibody (N-term) - Product Information

Application Primary Accession Other Accession

Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region FC, IHC-P, WB,E <u>P61313</u> <u>P61314</u>, <u>O9CZM2</u>, <u>O4R5B2</u>, <u>O5EAD6</u>, <u>NP_002939.2</u>, <u>G1T0C1</u> Human, Mouse Bovine, Monkey, Rabbit, Rat Rabbit Polyclonal Rabbit IgG 24146 21-50

RPL15 Antibody (N-term) - Additional Information

Gene ID 6138

Other Names 60S ribosomal protein L15, RPL15, EC45

Target/Specificity

This RPL15 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 21-50 amino acids from the N-terminal region of human RPL15.

Dilution FC~~1:10~50 IHC-P~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RPL15 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RPL15 Antibody (N-term) - Protein Information



Name RPL15

Synonyms EC45

Function Component of the large ribosomal subunit. The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell.


Cellular Location Cytoplasm.

RPL15 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

RPL15 Antibody (N-term) - Images




Anti-RPL15 Antibody (N-term) at 1:1000 dilution + HepG2 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





RPL15 Antibody (N-term) (Cat. #AP12772a)immunohistochemistry analysis in formalin fixed and paraffin embedded human pancreas tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of RPL15 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



RPL15 Antibody (N-term) (Cat. #AP12772a) flow cytometric analysis of K562 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated donkey-anti-rabbit secondary antibodies were used for the analysis.

RPL15 Antibody (N-term) - Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L15E family of ribosomal proteins. It is located in the cytoplasm. This gene shares sequence similarity with the yeast ribosomal protein YL10 gene. Although this gene has been referred to as RPL10, its official symbol is RPL15. This gene has been shown to be overexpressed in some esophageal tumors compared to normal matched tissues. Transcript variants utilizing alternative polyA signals exist. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

RPL15 Antibody (N-term) - References



Wheeler, H.E., et al. PLoS Genet. 5 (10), E1000685 (2009) : Wang, H., et al. BMC Cancer 6, 91 (2006) : Andersen, J.S., et al. Nature 433(7021):77-83(2005) Kapp, L.D., et al. Annu. Rev. Biochem. 73, 657-704 (2004) : Mazumder, B., et al. Cell 115(2):187-198(2003)