

## TRPC5 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP13769A

## Specification

# TRPC5 Antibody (N-term) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region IHC-P, WB,E <u>O9UL62</u> <u>O62852</u>, <u>O9OX29</u>, <u>NP\_036603.1</u>, <u>O9MYV9</u> Human Bovine, Mouse, Rabbit Rabbit Polyclonal Rabbit IgG 111412 254-283

# TRPC5 Antibody (N-term) - Additional Information

Gene ID 7224

**Other Names** Short transient receptor potential channel 5, TrpC5, Transient receptor protein 5, TRP-5, hTRP-5, hTRP5, TRPC5, TRPC5, TRP5

## Target/Specificity

This TRPC5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 254-283 amino acids from the N-terminal region of human TRPC5.

**Dilution** IHC-P~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

## Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

TRPC5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# **TRPC5 Antibody (N-term) - Protein Information**



Name TRPC5

Synonyms TRP5

**Function** Forms a receptor-activated non-selective calcium permeant cation channel (PubMed:<u>16284075</u>, PubMed:<u>38959890</u>, PubMed:<u>37137991</u>). Mediates calcium-dependent phosphatidylserine externalization and apoptosis in neurons via its association with PLSCR1 (By similarity). Acts on distinct neuronal populations in the hypothalamus to regulate innate behaviors including feeding, anxiety (flight/fight/fear), socialization, and maternal care (By similarity).

Cellular Location Cell membrane; Multi-pass membrane protein


**Tissue Location** Expressed in brain with higher levels in fetal brain. Found in cerebellum and occipital pole

# **TRPC5 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# TRPC5 Antibody (N-term) - Images



TRPC5 Antibody (N-term) (Cat. #AP13769a) western blot analysis in HL-60 cell line lysates (35ug/lane).This demonstrates the TRPC5 antibody detected the TRPC5 protein (arrow).





TRPC5 Antibody (N-term) (Cat. #AP13769a)immunohistochemistry analysis in formalin fixed and paraffin embedded human cerebellum tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of TRPC5 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.

# TRPC5 Antibody (N-term) - Background

This gene belongs to the transient receptor family. It encodes one of the seven mammalian TRPC (transient receptor potential channel) proteins. The encoded protein is a multi-pass membrane protein and is thought to form a receptor-activated non-selective calcium permeant cation channel. The protein is active alone or as a heteromultimeric assembly with TRPC1, TRPC3, and TRPC4. It also interacts with multiple proteins including calmodulin, CABP1, enkurin, Na(+)-H+ exchange regulatory factor (NHERF ), interferon-induced GTP-binding protein (MX1), ring finger protein 24 (RNF24), and SEC14 domain and spectrin repeat-containing protein 1 (SESTD1).

## **TRPC5 Antibody (N-term) - References**

Al-Shawaf, E., et al. Arterioscler. Thromb. Vasc. Biol. 30(7):1453-1459(2010) Wong, C.O., et al. Pflugers Arch. 460(1):121-130(2010) Miehe, S., et al. J. Biol. Chem. 285(16):12426-12434(2010) Gross, S.A., et al. J. Biol. Chem. 284(49):34423-34432(2009) Everett, K.V., et al. Hum. Genet. (2009) In press :