

CSTF3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14056b**Specification**

CSTF3 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q12996
Other Accession	Q99LI7 , NP_001317.1
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	82922
Antigen Region	556-585

CSTF3 Antibody (C-term) - Additional Information**Gene ID** 1479**Other Names**

Cleavage stimulation factor subunit 3, CF-1 77 kDa subunit, Cleavage stimulation factor 77 kDa subunit, CSTF 77 kDa subunit, CstF-77, CSTF3

Target/Specificity

This CSTF3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 556-585 amino acids from the C-terminal region of human CSTF3.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CSTF3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CSTF3 Antibody (C-term) - Protein Information**Name** CSTF3

Function One of the multiple factors required for polyadenylation and 3'-end cleavage of mammalian pre-mRNAs.

Cellular Location


Nucleus.

CSTF3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CSTF3 Antibody (C-term) - Images



CSTF3 Antibody (C-term) (Cat. #AP14056b) western blot analysis in ZR-75-1 cell line lysates (35ug/lane). This demonstrates the CSTF3 antibody detected the CSTF3 protein (arrow).

CSTF3 Antibody (C-term) - Background

The protein encoded by this gene is one of three (including CSTF1 and CSTF2) cleavage stimulation factors that combine to form the cleavage stimulation factor complex (CSTF). This complex is involved in the polyadenylation and 3' end cleavage of pre-mRNAs. The encoded protein functions as a homodimer and interacts directly with both CSTF1 and CSTF2 in the CSTF complex. Alternative splicing results in multiple transcript variants encoding different isoforms.

CSTF3 Antibody (C-term) - References

Hockert, J.A., et al. J. Biol. Chem. 285(1):695-704(2010)
Lamesch, P., et al. Genomics 89(3):307-315(2007)
Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :

Olsen, J.V., et al. Cell 127(3):635-648(2006)
Olsen, J.V., et al. Cell 127(3):635-648(2006)