

**SYT7 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP14342a**

**Specification**

---

**SYT7 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O43581</a>
Other Accession	<a href="#">NP_004191.2</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	45501
Antigen Region	72-101

**SYT7 Antibody (N-term) - Additional Information**

**Gene ID** 9066

**Other Names**

Synaptotagmin-7, IPCA-7, Prostate cancer-associated protein 7, Synaptotagmin VII, SytVII, SYT7, PCANAP7

**Target/Specificity**

This SYT7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 72-101 amino acids from the N-terminal region of human SYT7.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

SYT7 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**SYT7 Antibody (N-term) - Protein Information**

**Name** SYT7 ([HGNC:11514](#))

## Synonyms PCANAP7

**Function** Ca(2+) sensor involved in Ca(2+)-dependent exocytosis of secretory and synaptic vesicles through Ca(2+) and phospholipid binding to the C2 domain (By similarity). Ca(2+) induces binding of the C2- domains to phospholipid membranes and to assembled SNARE-complexes; both actions contribute to triggering exocytosis (By similarity). SYT7 binds Ca(2+) with high affinity and slow kinetics compared to other synaptotagmins (By similarity). Involved in Ca(2+)-triggered lysosomal exocytosis, a major component of the plasma membrane repair (PubMed:[11342594](#)). Ca(2+)-regulated delivery of lysosomal membranes to the cell surface is also involved in the phagocytic uptake of particles by macrophages (By similarity). Ca(2+)-triggered lysosomal exocytosis also plays a role in bone remodeling by regulating secretory pathways in osteoclasts and osteoblasts (By similarity). In case of infection, involved in participates cell invasion by Trypanosoma cruzi via Ca(2+)- triggered lysosomal exocytosis (PubMed:[11342594](#), PubMed:[15811535](#)). Involved in cholesterol transport from lysosome to peroxisome by promoting membrane contacts between lysosomes and peroxisomes: probably acts by promoting vesicle fusion by binding phosphatidylinositol-4,5- biphosphate on peroxisomal membranes (By similarity). Acts as a key mediator of synaptic facilitation, a process also named short-term synaptic potentiation: synaptic facilitation takes place at synapses with a low initial release probability and is caused by influx of Ca(2+) into the axon terminal after spike generation, increasing the release probability of neurotransmitters (By similarity). Probably mediates synaptic facilitation by directly increasing the probability of release (By similarity). May also contribute to synaptic facilitation by regulating synaptic vesicle replenishment, a process required to ensure that synaptic vesicles are ready for the arrival of the next action potential: SYT7 is required for synaptic vesicle replenishment by acting as a sensor for Ca(2+) and by forming a complex with calmodulin (By similarity). Also acts as a regulator of Ca(2+)- dependent insulin and glucagon secretion in beta-cells (By similarity). Triggers exocytosis by promoting fusion pore opening and fusion pore expansion in chromaffin cells (By similarity). Also regulates the secretion of some non-synaptic secretory granules of specialized cells (By similarity).

## Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q62747}; Single-pass membrane protein. Presynaptic cell membrane {ECO:0000250|UniProtKB:Q9R0N7}; Single-pass membrane protein. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane {ECO:0000250|UniProtKB:Q9R0N7}; Single-pass membrane protein. Lysosome membrane {ECO:0000250|UniProtKB:Q9R0N7}; Single-pass membrane protein. Cytoplasmic vesicle, phagosome membrane {ECO:0000250|UniProtKB:Q9R0N7}; Single-pass membrane protein. Peroxisome membrane {ECO:0000250|UniProtKB:Q9R0N7}; Single-pass membrane protein. Cytoplasmic vesicle, secretory vesicle membrane {ECO:0000250|UniProtKB:Q62747}; Single-pass membrane protein. Note=Localization to lysosomes is dependent on N- terminal palmitoylation and interaction with CD63 {ECO:0000250|UniProtKB:Q9R0N7}

## Tissue Location

Expressed in a variety of adult and fetal tissues.


## SYT7 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

## **SYT7 Antibody (N-term) - Images**



SYT7 Antibody (N-term) (Cat. #AP14342a) western blot analysis in human uterine tomer tissue lysates (35ug/lane).This demonstrates the SYT7 antibody detected the SYT7 protein (arrow).



SYT7 Antibody (N-term) (Cat. #AP14342a) western blot analysis in mouse heart tissue lysates (35ug/lane).This demonstrates the SYT7 antibody detected the SYT7 protein (arrow).

## **SYT7 Antibody (N-term) - Background**

Synaptotagmins, such as SYT7, are brain-specific calcium-dependent phospholipid-binding proteins that play a role in synaptic exocytosis and neurotransmitter release. See MIM 600782.

## **SYT7 Antibody (N-term) - References**

Bailey, S.D., et al. Diabetes Care (2010) In press :  
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)  
Fukuda, M., et al. Biochem. J. 365 (PT 1), 173-180 (2002) :  
Caler, E.V., et al. J. Exp. Med. 193(9):1097-1104(2001)  
Mizutani, A., et al. J. Biol. Chem. 275(13):9823-9831(2000)