

ALDH5A1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1472b**Specification**

ALDH5A1 Antibody (C-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	P51649
Other Accession	Q8BWF0
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	57215
Antigen Region	368-398

ALDH5A1 Antibody (C-term) - Additional Information**Gene ID** 7915**Other Names**

Succinate-semialdehyde dehydrogenase, mitochondrial, Aldehyde dehydrogenase family 5 member A1, NAD(+)-dependent succinic semialdehyde dehydrogenase, ALDH5A1, SSADH

Target/Specificity

This ALDH5A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 368-398 amino acids from the C-terminal region of human ALDH5A1.

Dilution

IHC-P~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ALDH5A1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ALDH5A1 Antibody (C-term) - Protein Information

Name ALDH5A1 ([HGNC:408](#))

Synonyms SSADH

Function Catalyzes one step in the degradation of the inhibitory neurotransmitter gamma-aminobutyric acid (GABA).

Cellular Location
Mitochondrion.


Tissue Location
Brain, pancreas, heart, liver, skeletal muscle and kidney. Lower in placenta

ALDH5A1 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ALDH5A1 Antibody (C-term) - Images



Western blot analysis of ALDH5A1 Antibody (C-term) (Cat.#AP1472b) in SK-BR-3 cell line lysates (35ug/lane). ALDH5A1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with *ALDH5A1 antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

ALDH5A1 Antibody (C-term) - Background

ALDH5A1 belongs to the aldehyde dehydrogenase family of proteins. This protein functions as a mitochondrial NAD(+)-dependent succinic semialdehyde dehydrogenase. A deficiency of this enzyme, known as 4-hydroxybutyricaciduria, is a rare inborn error in the metabolism of the neurotransmitter 4-aminobutyric acid (GABA). In response to the defect, physiologic fluids from patients accumulate GHB, a compound with numerous neuromodulatory properties.

ALDH5A1 Antibody (C-term) - References

Knerr,I., J. Inherit. Metab. Dis. 30 (3), 279-294 (2007)
Blasi,P., J. Mol. Evol. 63 (1), 54-68 (2006)
Plomin,R., Mol. Psychiatry 9 (6), 582-586 (2004)