

STC1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP14863b**Specification**

STC1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P52823
Other Accession	P97574 , O55183 , NP_003146.1
Reactivity	Human
Predicted	Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	27621
Antigen Region	193-221

STC1 Antibody (C-term) - Additional Information**Gene ID** 6781**Other Names**

Stanniocalcin-1, STC-1, STC1, STC

Target/Specificity

This STC1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 193-221 amino acids from the C-terminal region of human STC1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

STC1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

STC1 Antibody (C-term) - Protein Information**Name** STC1

Synonyms STC

Function Stimulates renal phosphate reabsorption, and could therefore prevent hypercalcemia.

Cellular Location
Secreted.

Tissue Location


Expressed in most tissues, with the highest levels in ovary, prostate, heart, kidney and thyroid. In the kidney, expression is confined to the nephron, specifically in the distal convoluted tubule and in the collecting tubule. Not detected in the brain, liver, spleen, peripheral blood leukocytes and adrenal medulla

STC1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STC1 Antibody (C-term) - Images



STC1 Antibody (C-term) (Cat. #AP14863b) western blot analysis in U251 cell line lysates (35ug/lane). This demonstrates the STC1 antibody detected the STC1 protein (arrow).

STC1 Antibody (C-term) - Background

This gene encodes a secreted, homodimeric glycoprotein that is expressed in a wide variety of tissues and may have autocrine or paracrine functions. The gene contains a 5' UTR rich in CAG trinucleotide repeats. The encoded protein contains 11 conserved cysteine residues and is phosphorylated by protein kinase C exclusively on its serine residues. The protein may play a role in the regulation of renal and intestinal calcium and phosphate

transport, cell metabolism, or cellular calcium/phosphate homeostasis. Overexpression of human stanniocalcin 1 in mice produces high serum phosphate levels, dwarfism, and increased metabolic rate. This gene has altered expression in hepatocellular, ovarian, and breast cancers.

STC1 Antibody (C-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Liu, G., et al. J. Natl. Cancer Inst. 102(11):812-827(2010)
Kottgen, A., et al. Nat. Genet. 42(5):376-384(2010)
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Trindade, D.M., et al. BMC Struct. Biol. 9, 57 (2009) :