

RPN1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16998b**Specification**

RPN1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P04843
Other Accession	P07153 , Q9GMB0 , Q4R4T0 , NP_002941.1
Reactivity	Human
Predicted	Monkey, Pig, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	68569
Antigen Region	559-587

RPN1 Antibody (C-term) - Additional Information**Gene ID** 6184**Other Names**

Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit 1,
Dolichyl-diphosphooligosaccharide--protein glycosyltransferase 67 kDa subunit, Ribophorin I,
RPN-I, Ribophorin-1, RPN1

Target/Specificity

This RPN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 559-587 amino acids from the C-terminal region of human RPN1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RPN1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RPN1 Antibody (C-term) - Protein Information

Name RPN1 ([HGNC:10381](#))

Function Subunit of the oligosaccharyl transferase (OST) complex that catalyzes the initial transfer of a defined glycan (Glc(3)Man(9)GlcNAc(2) in eukaryotes) from the lipid carrier dolichol-pyrophosphate to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains, the first step in protein N-glycosylation (PubMed:[31831667](#)). N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). All subunits are required for a maximal enzyme activity (By similarity).

Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:E2RQ08}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:E2RQ08}. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV.

Tissue Location


Expressed in all tissues tested.

RPN1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RPN1 Antibody (C-term) - Images



RPN1 Antibody (C-term) (Cat. #AP16998b) western blot analysis in WiDr cell line lysates (35ug/lane). This demonstrates the RPN1 antibody detected the RPN1 protein (arrow).

RPN1 Antibody (C-term) - Background

This gene encodes a type I integral membrane protein found only in the rough endoplasmic reticulum. The encoded protein is

part of an N-oligosaccharyl transferase complex that links high mannose oligosaccharides to asparagine residues found in the Asn-X-Ser/Thr consensus motif of nascent polypeptide chains. This protein forms part of the regulatory subunit of the 26S proteasome and may mediate binding of ubiquitin-like domains to this proteasome.

RPN1 Antibody (C-term) - References

Ruiz-Canada, C., et al. Cell 136(2):272-283(2009)
Wang, L., et al. Cancer Epidemiol. Biomarkers Prev. 17(12):3558-3566(2008)
Wilson, C.M., et al. Proc. Natl. Acad. Sci. U.S.A. 105(28):9534-9539(2008)
Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :
Chi, A., et al. J. Proteome Res. 5(11):3135-3144(2006)