

RCOR1 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17018A**Specification**

RCOR1 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	O9UKL0
Other Accession	O8CFE3 , NP_055971.1
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	53327
Antigen Region	101-130

RCOR1 Antibody (N-term) - Additional Information**Gene ID** 23186**Other Names**

REST corepressor 1, Protein CoREST, RCOR1, KIAA0071, RCOR

Target/Specificity

This RCOR1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 101-130 amino acids from the N-terminal region of human RCOR1.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RCOR1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RCOR1 Antibody (N-term) - Protein Information**Name** RCOR1

Synonyms KIAA0071, RCOR

Function Essential component of the BHC complex, a corepressor complex that represses transcription of neuron-specific genes in non-neuronal cells. The BHC complex is recruited at RE1/NRSE sites by REST and acts by deacetylating and demethylating specific sites on histones, thereby acting as a chromatin modifier. In the BHC complex, it serves as a molecular beacon for the recruitment of molecular machinery, including MeCP2 and SUV39H1, that imposes silencing across a chromosomal interval. Plays a central role in demethylation of Lys-4 of histone H3 by promoting demethylase activity of KDM1A on core histones and nucleosomal substrates. It also protects KDM1A from the proteasome. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development and controls hematopoietic differentiation.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00512, ECO:0000255|PROSITE-ProRule:PRU00624, ECO:0000269|PubMed:10734093, ECO:0000269|PubMed:15897453}. Note=Upon infection by HSV-1, it is partially translocated into the cytoplasm in an HSV-1-dependent manner


Tissue Location

Ubiquitously expressed.


RCOR1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RCOR1 Antibody (N-term) - Images

RCOR1 Antibody (N-term) (Cat. #AP17018a) western blot analysis in ZR-75-1 cell line lysates (35ug/lane). This demonstrates the RCOR1 antibody detected the RCOR1 protein (arrow).



All lanes : Anti-RCOR1 Antibody (N-term) at 1:2000 dilution Lane 1: HT-29 whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: LNCaP whole cell lysate Lane 4: MOLT-4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 53 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

RCOR1 Antibody (N-term) - Background

The RCOR gene encodes a functional corepressor required for regulation of neural-specific gene expression.[supplied by OMIM].

RCOR1 Antibody (N-term) - References

Xiao, J., et al. J. Orthop. Res. 28(12):1569-1575(2010)
Everett, R.D. J. Virol. 84(7):3695-3698(2010)
Das, C., et al. J. Mol. Biol. 397(1):1-12(2010)
Laurent, B., et al. Stem Cells 27(9):2153-2162(2009)
Wang, Y., et al. Cell 138(4):660-672(2009)