

ND4L Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17147b**Specification**

ND4L Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	P03901
Other Accession	Q7J3C3 , YP_003024034.1
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	65-93

ND4L Antibody (C-term) - Additional Information**Gene ID** 4539**Other Names**

NADH-ubiquinone oxidoreductase chain 4L, NADH dehydrogenase subunit 4L, MT-ND4L, MTND4L, NADH4L, ND4L

Target/Specificity

This ND4L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 65-93 amino acids from the C-terminal region of human ND4L.

Dilution

WB~~1:2000

FC~~1:25

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ND4L Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ND4L Antibody (C-term) - Protein Information**Name** MT-ND4L ([HGNC:7460](#))

Synonyms MTND4L, NADH4L, ND4L

Function Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor (PubMed:[28844695](#)). Part of the enzyme membrane arm which is embedded in the lipid bilayer and involved in proton translocation (PubMed:[28844695](#)).

Cellular Location


Mitochondrion inner membrane {ECO:0000250|UniProtKB:P03902}; Multi-pass membrane protein

ND4L Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ND4L Antibody (C-term) - Images



Anti-ND4L Antibody (C-term) at 1:2000 dilution + human skeletal muscle lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 10 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-ND4L Antibody (C-term) at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: HL-60 whole cell lysate Lane 3: Human kidney lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 11 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing U-2 OS cells stained with AP17147b(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP17147b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

ND4L Antibody (C-term) - Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).

ND4L Antibody (C-term) - References

Andrews, R.M., et al. Nat. Genet. 23 (2), 147 (1999) :
Anderson, S., et al. Nature 290(5806):457-465(1981)

Submitted (08-JUL-2009) National Center for Biotechnology Information, NIH, Bethesda, MD 20894, USA :

Kogelnik, A.M., et al. Submitted (24-AUG-2006) Mitomap.org, Center for Molecular and Mitochondrial Medicine and Genetics (MAMMAG) University of California, University of California, Irvine, Irvine, CA 92697-3940, USA :

Kogelnik, A.M., et al. Submitted (18-APR-1997) Center for Molecular Medicine, Emory University School of Medicine, 1462 Clifton Road, Suite 420, Atlanta, GA 30322, USA :