

PDGFB Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1721A

Specification

PDGFB Antibody (C-term) - Product Information

Application WB, IF, E **Primary Accession** P01127 Other Accession NP 148937 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 27283 Antigen Region 212-241

PDGFB Antibody (C-term) - Additional Information

Gene ID 5155

Other Names

Platelet-derived growth factor subunit B, PDGF subunit B, PDGF-2, Platelet-derived growth factor B chain, Platelet-derived growth factor beta polypeptide, Proto-oncogene c-Sis, Becaplermin, PDGFB, PDGF2, SIS

Target/Specificity

This PDGFB antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 212-241 amino acids from the C-terminal region of human PDGFB.

Dilution

WB~~1:1000 IF~~1:10~50

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PDGFB Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PDGFB Antibody (C-term) - Protein Information





Name PDGFB

Synonyms PDGF2, SIS

Function Growth factor that plays an essential role in the regulation of embryonic development, cell proliferation, cell migration, survival and chemotaxis. Potent mitogen for cells of mesenchymal origin (PubMed:26599395). Required for normal proliferation and recruitment of pericytes and vascular smooth muscle cells in the central nervous system, skin, lung, heart and placenta. Required for normal blood vessel development, and for normal development of kidney glomeruli. Plays an important role in wound healing. Signaling is modulated by the formation of heterodimers with PDGFA (By similarity).

Cellular Location

Secreted. Note=Released by platelets upon wounding

Tissue Location


Expressed at high levels in the heart, brain (sustantia nigra), placenta and fetal kidney. Expressed at moderate levels in the brain (hippocampus), skeletal muscle, kidney and lung

PDGFB Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

PDGFB Antibody (C-term) - Images



Confocal immunofluorescent analysis of PDGFB Antibody (C-term) (Cat#AP1721a) with 293 cell followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).





The anti-PDGFB C-term Pab (Cat. #AP1721a) is used in Western blot to detect PDGFB in A375 cell lysate.

PDGFB Antibody (C-term) - Background

PDGFB is a member of the platelet-derived growth factor family. The four members of this family are mitogenic factors for cells of mesenchymal origin and are characterized by a motif of eight cysteines. This gene product can exist either as a homodimer or as a heterodimer with the platelet-derived growth factor alpha polypeptide, where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Reciprocal translocations between chromosomes 22 and 7, at sites where this gene and that for COL1A1 are located, are associated with a particular type of skin tumor called dermatofibrosarcoma protuberans resulting from unregulated expression of growth factor.

PDGFB Antibody (C-term) - References

Muller, C., et al., J. Biol. Chem. 278(20):18330-18335 (2003). Laprise, M.H., et al., Blood 100(10):3578-3587 (2002). Maire, G., et al., Cancer Genet. Cytogenet. 134(2):156-161 (2002). Andrae, J., et al., Biochem. Biophys. Res. Commun. 296(3):604-611 (2002). Ehrlich, H.P., et al., Cytokines Cell Mol Ther 7(3):85-90 (2002).