

ARCN1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17651c**Specification**

ARCN1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P48444
Other Accession	Q66H80 , Q5XJY5 , P53619 , NP_001135753.1
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	57210
Antigen Region	296-324

ARCN1 Antibody (Center) - Additional Information**Gene ID** 372**Other Names**

Coatomer subunit delta, Archain, Delta-coat protein, Delta-COP, ARCN1, COPD

Target/Specificity

This ARCN1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 296-324 amino acids from the Central region of human ARCN1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ARCN1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ARCN1 Antibody (Center) - Protein Information**Name** ARCN1

Synonyms COPD

Function Component of the coatomer, a cytosolic protein complex that binds to dilysine motifs and reversibly associates with Golgi non- clathrin-coated vesicles, which further mediate biosynthetic protein transport from the ER, via the Golgi up to the trans Golgi network. The coatomer complex is required for budding from Golgi membranes, and is essential for the retrograde Golgi-to-ER transport of dilysine-tagged proteins. In mammals, the coatomer can only be recruited by membranes associated to ADP-ribosylation factors (ARFs), which are small GTP-binding proteins; the complex also influences the Golgi structural integrity, as well as the processing, activity, and endocytic recycling of LDL receptors (By similarity).

Cellular Location

Cytoplasm. Golgi apparatus membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, COPI-coated vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Note=The coatomer is cytoplasmic or polymerized on the cytoplasmic side of the Golgi, as well as on the vesicles/buds originating from it.

Tissue Location


Ubiquitously expressed.

ARCNI1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ARCNI1 Antibody (Center) - Images



ARCNI1 Antibody (Center) (Cat. #AP17651c) western blot analysis in ZR-75-1 cell line lysates (35ug/lane). This demonstrates the ARCNI1 antibody detected the ARCNI1 protein (arrow).

ARCNI1 Antibody (Center) - Background

This gene maps in a region, which include the mixed lineage leukemia and Friend leukemia virus integration 1 genes, where multiple disease-associated chromosome translocations occur. It is an intracellular protein. Archain sequences are well conserved among eukaryotes and this protein may play a fundamental role in eukaryotic cell biology. It has similarities to heat shock proteins and clathrin-associated proteins, and may be involved in vesicle structure or trafficking.

ARCNI1 Antibody (Center) - References

Lippincott-Schwartz, J., et al. Trends Cell Biol. 16 (10), E1-E4 (2006) :
Xu, Y., et al. Mol. Biol. Cell 13(10):3493-3507(2002)
Lippincott-Schwartz, J., et al. Annu. Rev. Cell Dev. Biol. 16, 557-589 (2000) :
Lowe, M., et al. J. Biol. Chem. 271(48):30725-30730(1996)
Tunnacliffe, A., et al. Mamm. Genome 7(10):784-786(1996)