

PNMA5 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17906B

Specification

PNMA5 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O96PV4
Other Accession	NP_001096620.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	49934
Antigen Region	387-416

PNMA5 Antibody (C-term) - Additional Information

Gene ID 114824

Other Names

Paraneoplastic antigen-like protein 5, Tumor antigen BJ-HCC-25, PNMA5, KIAA1934

Target/Specificity

This PNMA5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 387-416 amino acids from the C-terminal region of human PNMA5.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PNMA5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PNMA5 Antibody (C-term) - Protein Information

Name PNMA5

Synonyms KIAA1934

Cellular Location

Nucleus.


Tissue Location

Expressed in the brain.

PNMA5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PNMA5 Antibody (C-term) - Images

PNMA5 Antibody (C-term) (Cat. #AP17906b) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the PNMA5 antibody detected the PNMA5 protein (arrow).

PNMA5 Antibody (C-term) - Background

The function of this protein remains unknown.

PNMA5 Antibody (C-term) - References

Schuller, M., et al. J. Neuroimmunol. 169 (1-2), 172-176 (2005) :