

BARON Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1831c**Specification**

BARON Antibody (Center) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	Q92622
Other Accession	Q80U62
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	108622
Antigen Region	656-684

BARON Antibody (Center) - Additional Information**Gene ID** 9711**Other Names**

Run domain Beclin-1 interacting and cysteine-rich containing protein, Rubicon, Beclin-1 associated
RUN domain containing protein, Baron, KIAA0226

Target/Specificity

This BARON antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 656-684 amino acids from the Central region of human BARON.

Dilution

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BARON Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

BARON Antibody (Center) - Protein Information

Name RUBCN ([HGNC:28991](#))

Synonyms KIAA0226

Function Inhibits PI3K3 activity; under basal conditions negatively regulates PI3K complex II (PI3KC3-C2) function in autophagy. Negatively regulates endosome maturation and degradative endocytic trafficking and impairs autophagosome maturation process. Can sequester UVRAG from association with a class C Vps complex (possibly the HOPS complex) and negatively regulates Rab7 activation (PubMed:[20974968](#), PubMed:[21062745](#)).

Cellular Location


Late endosome. Lysosome. Early endosome Note=Predominantly located in late endosomes/lysosomes, only partially detected in early endosome and not at all in the Golgi apparatus

BARON Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


BARON Antibody (Center) - Images



All lanes : Anti-BARON Antibody (Center) at 1:1000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: Raji whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 109 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human kidney reacted with BARON Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



BARON Antibody (Center) (Cat. #AP1831c) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

BARON Antibody (Center) - References

Matsunaga, K., et al. Nat. Cell Biol. 11(4):385-396(2009)
Muzny, D.M., et al. Nature 440(7088):1194-1198(2006)
Venter, J.C., et al. Science 291(5507):1304-1351(2001)