

MC5R Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18970b

Specification

MC5R Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P33032
Other Accession	P56451 , B0V1P1 , NP_005904.1 , P41983
Reactivity	Human
Predicted	Zebrafish, Bovine, Sheep
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36601
Antigen Region	280-308

MC5R Antibody (C-term) - Additional Information

Gene ID 4161

Other Names

Melanocortin receptor 5, MC5-R, MC-2, MC5R

Target/Specificity

This MC5R antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 280-308 amino acids from the C-terminal region of human MC5R.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MC5R Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MC5R Antibody (C-term) - Protein Information

Name MC5R

Function Receptor for MSH (alpha, beta and gamma) and ACTH. The activity of this receptor is mediated by G proteins which activate adenylate cyclase. This receptor is a possible mediator of the immunomodulation properties of melanocortins.

Cellular Location

Cell membrane; Multi-pass membrane protein.

Tissue Location


Expressed in the brain but not in the melanoma cells

MC5R Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MC5R Antibody (C-term) - Images



MC5R Antibody (C-term) (Cat. #AP18970b) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the MC5R antibody detected the MC5R protein (arrow).



Anti-MC5R Antibody (C-term) at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

MC5R Antibody (C-term) - Background

This gene encodes a member of the seven-pass transmembrane G protein-coupled melanocortin receptor protein family that stimulate cAMP signal transduction. The encoded protein is a receptor for melanocyte-stimulating hormone and adrenocorticotrophic hormone and is suggested to play a role in sebum generation.

MC5R Antibody (C-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Pinheiro, A.P., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (5), 1070-1080 (2010) :
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Miller, C.L., et al. Schizophr. Res. 113 (2-3), 259-267 (2009) :
Frandsberg, P.A., et al. Biochem. Biophys. Res. Commun. 236(2):489-492(1997)