

**EIF3E Antibody(Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP19409c****Specification**

---

**EIF3E Antibody(Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P60228</a>
Other Accession	<a href="#">Q641X8</a> , <a href="#">P60229</a> , <a href="#">Q4R6G8</a> , <a href="#">Q5ZLA5</a> , <a href="#">Q3T102</a> , <a href="#">Q3B8M3</a> , <a href="#">Q1LUA8</a> , <a href="#">Q05AY2</a> , <a href="#">Q6DRI1</a> , <a href="#">NP_001559.1</a>
Reactivity	Human, Zebrafish
Predicted	Xenopus, Bovine, Chicken, Monkey, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52221
Antigen Region	248-276

**EIF3E Antibody(Center) - Additional Information****Gene ID** 3646**Other Names**

Eukaryotic translation initiation factor 3 subunit E {ECO:0000255|HAMAP-Rule:MF\_03004}, eIF3e {ECO:0000255|HAMAP-Rule:MF\_03004}, Eukaryotic translation initiation factor 3 subunit 6 {ECO:0000255|HAMAP-Rule:MF\_03004}, Viral integration site protein INT-6 homolog, eIF-3 p48 {ECO:0000255|HAMAP-Rule:MF\_03004}, EIF3E {ECO:0000255|HAMAP-Rule:MF\_03004}

**Target/Specificity**

This EIF3E antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 248-276 amino acids from the Central region of human EIF3E.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

EIF3E Antibody(Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## EIF3E Antibody(Center) - Protein Information

**Name** EIF3E {ECO:0000255|HAMAP-Rule:MF\_03004}

**Function** Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed:[17581632](#), PubMed:[25849773](#), PubMed:[27462815](#)). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNA<sup>i</sup> and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed:[17581632](#)). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed:[25849773](#)). Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway (PubMed:[17468741](#)). May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins (PubMed:[17310990](#), PubMed:[17324924](#)).

### Cellular Location

Cytoplasm. Nucleus, PML body.

### Tissue Location


Ubiquitously expressed. Expressed at highest levels in appendix, lymph, pancreas, skeletal muscle, spleen and thymus


## EIF3E Antibody(Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## EIF3E Antibody(Center) - Images





EIF3E Antibody (Center)(Cat. #AP19409c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the EIF3E antibody detected the EIF3E protein (arrow).



(DANRE) eif3eb Antibody (Center) (Cat. # AP19409c) western blot analysis in zebra fish brain tissue lysates (35ug/lane). This demonstrates the (DANRE) eif3eb antibody detected the (DANRE) eif3eb protein (arrow).

### EIF3E Antibody(Center) - Background

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA<sub>i</sub> and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway. May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins.

### EIF3E Antibody(Center) - References

Grzmil, M., et al. *Oncogene* 29(28):4080-4089(2010)  
Zhou, M., et al. *Proc. Natl. Acad. Sci. U.S.A.* 105(47):18139-18144(2008)  
Masutani, M., et al. *EMBO J.* 26(14):3373-3383(2007)

Morris, C., et al. EMBO Rep. 8(6):596-602(2007)  
Sirchia, R., et al. Biol. Chem. 388(5):457-465(2007)