

### **GAPT Antibody(Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP19800C

### Specification

# **GAPT Antibody(Center) - Product Information**

Application	WB,E
Primary Accession	<u>Q8N292</u>
Other Accession	<u>NP_689900.1</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	17883
Antigen Region	84-112

### GAPT Antibody(Center) - Additional Information

### Gene ID 202309

### Other Names

Protein GAPT, Growth factor receptor-bound protein 2-binding adapter protein, transmembrane, GAPT, C5orf29

#### Target/Specificity

This GAPT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 84-112 amino acids from the Central region of human GAPT.

Dilution WB~~1:1000

 $E \sim Use$  at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

GAPT Antibody(Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# **GAPT Antibody(Center) - Protein Information**

Name GAPT



# Synonyms C5orf29

**Function** Negatively regulates B-cell proliferation following stimulation through the B-cell receptor. May play an important role in maintenance of marginal zone (MZ) B-cells (By similarity).

**Cellular Location** Cell membrane; Single-pass membrane protein

Tissue Location


Highly expressed in spleen and PBL, detected at lower levels in thymus, and undetectable in all other tissues tested Also expressed in various B-cell lines, monocytic cell line THP-1 and NK-like cell line YT, but not in T-cell line Jurkat or HeLa cells

# GAPT Antibody(Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# GAPT Antibody(Center) - Images



GAPT Antibody (Center) (Cat. #AP19800c) western blot analysis in NCI-H460 cell line lysates (35ug/lane).This demonstrates the GAPT antibody detected the GAPT protein (arrow).

# GAPT Antibody(Center) - Background

Negatively regulates B-cell proliferation following stimulation through the B-cell receptor. May play an important role in maintenance of marginal zone (MZ) B-cells (By similarity).

# GAPT Antibody(Center) - References

Gerhard, D.S., et al. Genome Res. 14 (10B), 2121-2127 (2004) :