

**APOBEC3A Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20219a****Specification**

---

**APOBEC3A Antibody (N-term) - Product Information**

|                   |                             |
|-------------------|-----------------------------|
| Application       | WB,E                        |
| Primary Accession | <a href="#">P31941</a>      |
| Other Accession   | <a href="#">NP_663745.1</a> |
| Reactivity        | Human                       |
| Host              | Rabbit                      |
| Clonality         | Polyclonal                  |
| Isotype           | Rabbit IgG                  |
| Calculated MW     | 23012                       |
| Antigen Region    | 20-49                       |

**APOBEC3A Antibody (N-term) - Additional Information****Gene ID** 100913187;200315**Other Names**

DNA dC-&gt;dU-editing enzyme APOBEC-3A, A3A, 354-, Phorbolin-1, APOBEC3A

**Target/Specificity**

This APOBEC3A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 20-49 amino acids from the N-terminal region of human APOBEC3A.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

APOBEC3A Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**APOBEC3A Antibody (N-term) - Protein Information****Name** APOBEC3A**Function** DNA deaminase (cytidine deaminase) with restriction activity against viruses, foreign

DNA and mobility of retrotransposons. Exhibits antiviral activity against adeno-associated virus (AAV) and human T- cell leukemia virus type 1 (HTLV-1) and may inhibit the mobility of LTR and non-LTR retrotransposons. Selectively targets single-stranded DNA and can deaminate both methylcytosine and cytosine in foreign DNA. Can induce somatic hypermutation in the nuclear and mitochondrial DNA. May also play a role in the epigenetic regulation of gene expression through the process of active DNA demethylation.

**Cellular Location**

Nucleus. Cytoplasm.


**Tissue Location**

Expressed in peripheral leukocytes with higher expression in CD14-positive phagocytic cells. Highly expressed in keratinocytes and in periphery blood monocytes. Also detected in non-lymphoid tissues including lung and adipose tissues. Found at high levels in colorectal adenocarcinoma, Burkitt's lymphoma and chronic myelogenous leukemia.

**APOBEC3A Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**APOBEC3A Antibody (N-term) - Images**

Anti-APOBEC3A Antibody (N-term) at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

**APOBEC3A Antibody (N-term) - Background**

This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. The protein encoded by this gene lacks the zinc binding activity of other family members. The protein plays a role in immunity, by restricting transmission of foreign DNA such as viruses. One mechanism of foreign DNA restriction is deamination of foreign double-stranded DNA cytidines to uridines, which leads to DNA degradation. However, other mechanisms are also thought to be involved, as anti-viral effect is not dependent on deaminase activity. One allele of this gene results from the deletion of approximately 29.5 kb of sequence between this gene, APOBEC3A, and the adjacent gene APOBEC3B. The breakpoints of the deletion are within the two genes, so the deletion allele is predicted to have the promoter and coding region of APOBEC3A, but the 3' UTR of APOBEC3B.

#### **APOBEC3A Antibody (N-term) - References**

Thielen, B.K., et al. J. Biol. Chem. 285(36):27753-27766(2010)  
Berger, A., et al. J. Biol. Chem. 285(16):12248-12254(2010)  
Stenglein, M.D., et al. Nat. Struct. Mol. Biol. 17(2):222-229(2010)  
Abe, H., et al. Hepatol. Res. 39(12):1159-1168(2009)  
Prochnow, C., et al. Sci. China, C, Life Sci. 52(10):893-902(2009)

#### **APOBEC3A Antibody (N-term) - Citations**

- [Comprehensive Analyses Identify APOBEC3A as a Genomic Instability-Associated Immune Prognostic Biomarker in Ovarian Cancer](#)