

ESD Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20553c**Specification**

ESD Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P10768
Other Accession	O9GJT2
Reactivity	Human, Mouse, Rat
Predicted	Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	31463

ESD Antibody (Center) - Additional Information**Gene ID** 2098**Other Names**

S-formylglutathione hydrolase, FGH, Esterase D, Methylumbelliferyl-acetate deacetylase, ESD

Target/Specificity

This ESD antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 68-102 amino acids from the Central region of human ESD.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ESD Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ESD Antibody (Center) - Protein Information**Name** ESD**Function** Serine hydrolase involved in the detoxification of formaldehyde.


Cellular Location

Cytoplasm. Cytoplasmic vesicle.

ESD Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ESD Antibody (Center) - Images

Western blot analysis of lysates from Jurkat, K562 cell line, mouse stomach and rat stomach tissue (from left to right), using ESD Antibody (Center) (Cat. #AP20553c). AP20553c was diluted at 1:1000 at each lane. A goat anti-rabbit(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

ESD Antibody (Center) - Background

Serine hydrolase involved in the detoxification of formaldehyde.

ESD Antibody (Center) - References

Lee E.Y.-H.P.,et al.Proc. Natl. Acad. Sci. U.S.A. 83:6337-6341(1986).
Young L.-J.S.,et al.Hum. Genet. 79:137-141(1988).
Hu R.-M.,et al.Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Dunham A.,et al.Nature 428:522-528(2004).