

DFFB Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20612c

Specification

DFFB Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	O76075 , O54788
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG

DFFB Antibody (Center) - Additional Information

Other Names

DNA fragmentation factor subunit beta, 3---, Caspase-activated deoxyribonuclease, CAD, Caspase-activated DNase, Caspase-activated nuclease, CPAN, DNA fragmentation factor 40 kDa subunit, DFF-40, DFFB, CAD, DFF2, DFF40

Target/Specificity

This DFFB antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 203-238 amino acids from the Central region of human DFFB.

Dilution

WB~~1:1000
IHC-P~~1:25
FC~~1:25
E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DFFB Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.


DFFB Antibody (Center) - Protein Information

DFFB Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


DFFB Antibody (Center) - Images




Western blot analysis of lysate from A431 cell line, using DFFB Antibody (Center) (Cat. #AP20612c). AP20612c was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Immunohistochemical analysis of paraffin-embedded mouse kidney section using DFFB Antibody (Center) (Cat.#AP20612c). AP20612c was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded human kidney section using DFFB Antibody (Center) (Cat#AP20612c). AP20612c was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Flow cytometric analysis of Hela cells using DFFB Antibody (Center) (green, Cat#AP20612c) compared to an isotype control of rabbit IgG (blue). AP20612c was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

DFFB Antibody (Center) - Background

Nuclease that induces DNA fragmentation and chromatin condensation during apoptosis.
Degrades naked DNA and induces apoptotic morphology.

DFFB Antibody (Center) - References

Liu X.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:8461-8466(1998).
Mukae N.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:9123-9128(1998).
Halenbeck R.,et al.Curr. Biol. 8:537-540(1998).
Nakagawara A.,et al.Submitted (JUN-1999) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).