

DHFR Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20626a**Specification**

DHFR Antibody (N-term) - Product Information

Application	IHC-P, WB,E
Primary Accession	P00374
Other Accession	P00377 , P00375 , P00376
Reactivity	Human, Mouse, Rat
Predicted	Bovine, Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21453

DHFR Antibody (N-term) - Additional Information**Gene ID** 1719**Other Names**

Dihydrofolate reductase, DHFR

Target/Specificity

This DHFR antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 16-30 amino acids from the N-terminal region of human DHFR.

Dilution

IHC-P~~1:25

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

DHFR Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

DHFR Antibody (N-term) - Protein Information**Name** DHFR

Function Key enzyme in folate metabolism. Contributes to the de novo mitochondrial thymidylate biosynthesis pathway. Catalyzes an essential reaction for de novo glycine and purine synthesis, and for DNA precursor synthesis. Binds its own mRNA and that of DHFR2.

Cellular Location

Mitochondrion {ECO:0000250|UniProtKB:P00375}. Cytoplasm {ECO:0000250|UniProtKB:P00375}

Tissue Location


Widely expressed in fetal and adult tissues, including throughout the fetal and adult brains and whole blood Expression is higher in the adult brain than in the fetal brain

DHFR Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


DHFR Antibody (N-term) - Images



Western blot analysis of lysates from 293T, HeLa, Jurkat, Y79 cell line, mouse spleen and rat kidney tissue lysate (from left to right), using DHFR Antibody (N-term) (Cat. #AP20626a). AP20626a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Immunohistochemical analysis of paraffin-embedded H. liver section using DHFR Antibody (N-term)(Cat#AP20626a). AP20626a was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. brain section using DHFR Antibody (N-term)(Cat#AP20626a). AP20626a was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

DHFR Antibody (N-term) - Background

Key enzyme in folate metabolism. Contributes to the de novo mitochondrial thymidylate biosynthesis pathway. Catalyzes an essential reaction for de novo glycine and purine synthesis, and for DNA precursor synthesis. Binds its own mRNA and that of DHFRL1.

DHFR Antibody (N-term) - References

Chen M.-J.,et al.J. Biol. Chem. 259:3933-3943(1984).
Masters J.N.,et al.Gene 21:59-63(1983).
Yang J.K.,et al.J. Mol. Biol. 176:169-187(1984).
Schmutz J.,et al.Nature 431:268-274(2004).
Banka S.,et al.Am. J. Hum. Genet. 88:216-225(2011).