

ECHDC3 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20684a**Specification**

ECHDC3 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q96DC8
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG

ECHDC3 Antibody (N-term) - Additional Information**Gene ID** 79746**Other Names**

Enoyl-CoA hydratase domain-containing protein 3, mitochondrial, ECHDC3

Target/Specificity

This ECHDC3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 31-66 amino acids from the N-terminal region of human ECHDC3.

Dilution

WB~~1:1000

IHC-P~~1:25

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ECHDC3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ECHDC3 Antibody (N-term) - Protein Information**Name** ECHDC3 ([HGNC:23489](#))**Function** May play a role in fatty acid biosynthesis and insulin sensitivity.**Cellular Location**

Mitochondrion.

Tissue Location


Expressed in adipocytes (PubMed:31010960). Expressed in blood cells, with higher expression in patients with low coronary lesions (PubMed:27586541).

ECHDC3 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ECHDC3 Antibody (N-term) - Images



Western blot analysis of lysates from human liver and kidney tissue lysate (from left to right), using ECHDC3 Antibody (N-term)(Cat. #AP20684a). AP20684a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Immunohistochemical analysis of paraffin-embedded H. kidney section using ECHDC3 Antibody (N-term)(Cat#AP20684a). AP20684a was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

ECHDC3 Antibody (N-term) - References

- Ota T.,et al.Nat. Genet. 36:40-45(2004).
Wan D.,et al.Proc. Natl. Acad. Sci. U.S.A. 101:15724-15729(2004).
Suzuki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Deloukas P.,et al.Nature 429:375-381(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.