

BAX Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20807a**Specification**

BAX Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q07812
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	21184

BAX Antibody (N-term) - Additional Information**Gene ID** 581**Other Names**

Apoptosis regulator BAX, Bcl-2-like protein 4, Bcl2-L-4, BAX, BCL2L4

Target/Specificity

This BAX antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 47-78 amino acids from the N-terminal region of human BAX.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BAX Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

BAX Antibody (N-term) - Protein Information**Name** BAX**Synonyms** BCL2L4**Function** Plays a role in the mitochondrial apoptotic process (PubMed:[10772918](#),

PubMed:[11060313](#), PubMed:[16113678](#), PubMed:[16199525](#), PubMed:[18948948](#), PubMed:[21199865](#), PubMed:[21458670](#), PubMed:[25609812](#), PubMed:[36361894](#), PubMed:[8358790](#), PubMed:[8521816](#)). Under normal conditions, BAX is largely cytosolic via constant retrotranslocation from mitochondria to the cytosol mediated by BCL2L1/Bcl-xL, which avoids accumulation of toxic BAX levels at the mitochondrial outer membrane (MOM) (PubMed:[21458670](#)). Under stress conditions, undergoes a conformation change that causes translocation to the mitochondrion membrane, leading to the release of cytochrome c that then triggers apoptosis (PubMed:[10772918](#), PubMed:[11060313](#), PubMed:[16113678](#), PubMed:[16199525](#), PubMed:[18948948](#), PubMed:[21199865](#), PubMed:[21458670](#), PubMed:[25609812](#), PubMed:[8358790](#), PubMed:[8521816](#)). Promotes activation of CASP3, and thereby apoptosis (PubMed:[10772918](#), PubMed:[11060313](#), PubMed:[16113678](#), PubMed:[16199525](#), PubMed:[18948948](#), PubMed:[21199865](#), PubMed:[21458670](#), PubMed:[25609812](#), PubMed:[8358790](#), PubMed:[8521816](#)).

Cellular Location

[Isoform Alpha]: Mitochondrion outer membrane; Single-pass membrane protein. Cytoplasm. Nucleus Note=Colocalizes with 14-3-3 proteins in the cytoplasm. Under stress conditions, undergoes a conformation change that causes release from JNK-phosphorylated 14-3-3 proteins and translocation to the mitochondrion membrane. Upon Sendai virus infection, recruited to the mitochondrion through interaction with IRF3 (PubMed:[25609812](#)) [Isoform Gamma]: Cytoplasm.

Tissue Location


Expressed in a wide variety of tissues. Isoform Psi is found in glial tumors. Isoform Alpha is expressed in spleen, breast, ovary, testis, colon and brain, and at low levels in skin and lung. Isoform Sigma is expressed in spleen, breast, ovary, testis, lung, colon, brain and at low levels in skin. Isoform Alpha and isoform Sigma are expressed in pro-myelocytic leukemia, histiocytic lymphoma, Burkitt's lymphoma, T-cell lymphoma, lymphoblastic leukemia, breast adenocarcinoma, ovary adenocarcinoma, prostate carcinoma, prostate adenocarcinoma, lung carcinoma, epidermoid carcinoma, small cell lung carcinoma and colon adenocarcinoma cell lines


BAX Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BAX Antibody (N-term) - Images





Western blot analysis of lysates from HeLa, 293, HT-1080, Raji, SH-SY5Y, THP-1 cell line (from left to right), using BAX Antibody (N-term)(Cat. #AP20807a). AP20807a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35ug per lane.



Anti-BAX Antibody (N-term) at 1:2000 dilution + U-2OS whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

BAX Antibody (N-term) - Background

Accelerates programmed cell death by binding to, and antagonizing the apoptosis repressor BCL2 or its adenovirus homolog E1B 19k protein. Under stress conditions, undergoes a conformation change that causes translocation to the mitochondrion membrane, leading to the release of cytochrome c that then triggers apoptosis. Promotes activation of CASP3, and thereby apoptosis.

BAX Antibody (N-term) - References

- Oltvai Z.N.,et al.Cell 74:609-619(1993).
- Apte S.S.,et al.Genomics 26:592-594(1995).
- Shi B.,et al.Biochem. Biophys. Res. Commun. 254:779-785(1999).
- Schmitt E.,et al.Biochem. Biophys. Res. Commun. 270:868-879(2000).

Cartron P.F.,et al.Hum. Mol. Genet. 11:675-687(2002).