

**(Mouse) Dppa3 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP20950c**

**Specification**

---

**(Mouse) Dppa3 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8QZY3</a>
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	17670

**(Mouse) Dppa3 Antibody (C-term) - Additional Information**

**Gene ID** 73708

**Other Names**

Developmental pluripotency-associated protein 3, Compaction-associated protein 1, Primordial germ cell protein 7, Dppa3, Cap1p, Crg1, Pgc7

**Target/Specificity**

This mouse Dppa3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 134-170 amino acids from the C-terminal region of mouse Dppa3.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

(Mouse) Dppa3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**(Mouse) Dppa3 Antibody (C-term) - Protein Information**

**Name** Dppa3

**Synonyms** Cap1p, Crg1, Pgc7 {ECO:0000303|PubMed:11

**Function** Primordial germ cell (PGCs)-specific protein involved in epigenetic chromatin reprogramming in the zygote following fertilization. In zygotes, DNA demethylation occurs selectively in the paternal pronucleus before the first cell division, while the adjacent maternal pronucleus and certain paternally-imprinted loci are protected from this process. Participates in protection of DNA methylation in the maternal pronucleus by preventing conversion of 5mC to 5hmC: specifically recognizes and binds histone H3 dimethylated at 'Lys-9' (H3K9me2) on maternal genome, and protects maternal genome from TET3- mediated conversion to 5hmC and subsequent DNA demethylation. Does not bind paternal chromatin, which is mainly packed into protamine and does not contain much H3K9me2 mark. Also protects imprinted loci that are marked with H3K9me2 in mature sperm from DNA demethylation in early embryogenesis. May be important for the totipotent/pluripotent states continuing through preimplantation development. Also involved in chromatin condensation in oocytogenesis.

#### **Cellular Location**

Nucleus. Cytoplasm Note=Localized in the cytoplasm at the primary oocyte stage and in oocytes within mono-laminar follicles. Expressed in the nucleus and cytoplasm of oocytes in bi-laminar and Graafian follicles and during the 2-cell and morula stages. In 3.5 dpc blastocysts localization is mainly nuclear. Mainly localizes in the female pronucleus, localization to the male pronucleus is much weaker.

#### **Tissue Location**

Expressed in the immature oocytes and in newborn ovaries. Subsequently detected in maturing oocytes and in preimplantation embryos. Expressed in pluripotent embryonic but not in differentiated somatic cells. Expressed in blastocysts, epiblasts, primordial germ cells, embryonic gonads and primitive spermatogonia. No expression is detected in adult testes.


#### **(Mouse) Dppa3 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **(Mouse) Dppa3 Antibody (C-term) - Images**





Western blot analysis of lysates from mouse liver, mouse testis tissue (from left to right), using (Mouse) Dppa3 Antibody (C-term)(Cat. #AP20950c). AP20950c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.

#### **(Mouse) Dppa3 Antibody (C-term) - Background**

Primordial germ cell (PGCs)-specific protein involved in epigenetic chromatin reprogramming in the zygote following fertilization. In zygotes, DNA demethylation occurs selectively in the paternal pronucleus before the first cell division, while the adjacent maternal pronucleus and certain paternally-imprinted loci are protected from this process. Participates in protection of DNA methylation in the maternal pronucleus by preventing conversion of 5mC to 5hmC: specifically recognizes and binds histone H3 dimethylated at 'Lys-9' (H3K9me2) on maternal genome, and protects maternal genome from TET3-mediated conversion to 5hmC and subsequent DNA demethylation. Does not bind paternal chromatin, which is mainly packed into protamine and does not contain much H3K9me2 mark. Also protects imprinted loci that are marked with H3K9me2 in mature sperm from DNA demethylation in early embryogenesis. May be important for the totipotent/pluripotent states continuing through preimplantation development. Also involved in chromatin condensation in oocytogenesis.

#### **(Mouse) Dppa3 Antibody (C-term) - References**

Saitou M.,et al.Nature 418:293-300(2002).  
Sato M.,et al.Mech. Dev. 113:91-94(2002).  
Bortvin A.,et al.Development 130:1673-1680(2003).  
Li W.,et al.Submitted (JUL-2002) to the EMBL/GenBank/DDBJ databases.  
Carninci P.,et al.Science 309:1559-1563(2005).