

NOVA1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20963a

Specification

NOVA1 Antibody (Center) - Product Information

Application Primary Accession Other Accession	IHC-P, WB,E <u>P51513</u> <u>O9IKN6</u>
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	51727

NOVA1 Antibody (Center) - Additional Information

Gene ID 4857

Other Names

RNA-binding protein Nova-1, Neuro-oncological ventral antigen 1, Onconeural ventral antigen 1, Paraneoplastic Ri antigen, Ventral neuron-specific protein 1, NOVA1

Target/Specificity

This NOVA1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 139-173 amino acids from the Central region of human NOVA1.

Dilution IHC-P~~1:25 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NOVA1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NOVA1 Antibody (Center) - Protein Information

Name NOVA1 (HGNC:7886)



Function Functions to regulate alternative splicing in neurons by binding pre-mRNA in a sequence-specific manner to activate exon inclusion or exclusion. It binds specifically to the sequences 5'-YCAY- 3' and regulates splicing in only a subset of regulated exons (PubMed:<u>10811881</u>). Binding to an exonic 5'-YCAY-3' cluster changes the protein complexes assembled on pre-mRNA, blocking U1 snRNP binding and exon inclusion, whereas binding to an intronic 5'-YCAY-3' cluster enhances spliceosome assembly and exon inclusion. Binding to 5'-YCAY-3' clusters results in a local and asymmetric action to regulate spliceosome assembly and alternative splicing in neurons. Binding to an exonic 5'-YCAY-3' cluster changed the protein complexes assembled on pre-mRNA, blocking U1 snRNP (small nuclear ribonucleoprotein) binding and exon inclusion, whereas binding to an intronic 5'-YCAY-3' cluster enhanced spliceosome assembly and exon inclusion. With NOVA1, they perform unique biological functions in different brain areas and cell types. Autoregulates its own expression by acting as a splicing repressor. Acts to activate the inclusion of exon E3A in the glycine receptor alpha-2 chain and of exon E9 in gamma-aminobutyric-acid receptor gamma-2 subunit via a distal downstream UCAU-rich intronic splicing enhancer. Acts to regulate a novel glycine receptor alpha-2 chain splice variant (alpha-2N) in developing spinal cord (By similarity).

Cellular Location Nucleus {ECO:0000250|UniProtKB:Q9JKN6}.


Tissue Location Expressed in cerebellum, brain stem, hippocampus, and frontal cortex.

NOVA1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

NOVA1 Antibody (Center) - Images





Western blot analysis of lysates from A431, Hela cell line (from left to right), using NOVA1 Antibody (Center)(Cat. #AP20963a). AP20963a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20ug per lane.



Immunohistochemical analysis of paraffin-embedded H. astroglioma section using NOVA1 Antibody (Center)(Cat#AP20963a). AP20963a was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

NOVA1 Antibody (Center) - Background

May regulate RNA splicing or metabolism in a specific subset of developing neurons.

NOVA1 Antibody (Center) - References

Buckanovich R.J., et al.Neuron 11:657-672(1993). Ota T., et al.Nat. Genet. 36:40-45(2004). Venter J.C., et al.Science 291:1304-1351(2001). Mural R.J., et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Dmitrenko V.V., et al.Submitted (APR-1996) to the EMBL/GenBank/DDBJ databases.