

Bmi1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21436b**Specification**

Bmi1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P35226
Other Accession	P25916
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG

Bmi1 Antibody (C-term) - Additional Information**Gene ID** 100532731;648**Other Names**

Polycomb complex protein BMI-1, Polycomb group RING finger protein 4, Bmi1, Bmi-1, Pcgf4

Target/Specificity

This Bmi1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 235-268 amino acids from the C-terminal region of human Bmi1.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Bmi1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Bmi1 Antibody (C-term) - Protein Information**Name** BMI1**Synonyms** PCGF4, RNF51

Function Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:[15386022](#), PubMed:[16359901](#), PubMed:[16714294](#), PubMed:[21772249](#), PubMed:[25355358](#), PubMed:[26151332](#), PubMed:[27827373](#)). The complex composed of RNF2, UB2D3 and BMI1 binds nucleosomes, and has activity only with nucleosomal histone H2A (PubMed:[21772249](#), PubMed:[25355358](#)). In the PRC1-like complex, regulates the E3 ubiquitin-protein ligase activity of RNF2/RING2 (PubMed:[15386022](#), PubMed:[21772249](#), PubMed:[26151332](#)).

Cellular Location


Nucleus. Cytoplasm

Bmi1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Bmi1 Antibody (C-term) - Images



Anti-Bmi1 Antibody (C-term) at 1:2000 dilution + H-4-II-E whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Bmi1 Antibody (C-term) - Background

Component of a Polycomb group (PcG) multiprotein PRC1- like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout

development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. In the PRC1 complex, it is required to stimulate the E3 ubiquitin-protein ligase activity of RNF2/RING2 (By similarity).

Bmi1 Antibody (C-term) - References

Haupt Y.,et al.Cell 65:753-763(1991).
van Lohuiizen M.,et al.Cell 65:737-752(1991).
Cao R.,et al.Mol. Cell 20:845-854(2005).
Buchwald G.,et al.EMBO J. 25:2465-2474(2006).