

**CNP Antibody (N-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21577a****Specification**

---

**CNP Antibody (N-Term) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB,E                   |
| Primary Accession | <a href="#">P09543</a> |
| Reactivity        | Mouse, Rat             |
| Host              | Rabbit                 |
| Clonality         | polyclonal             |
| Isotype           | Rabbit IgG             |
| Calculated MW     | 47579                  |

**CNP Antibody (N-Term) - Additional Information****Gene ID** 1267**Other Names**

2', 3'-cyclic-nucleotide 3'-phosphodiesterase, CNP, CNPase, CNP

**Target/Specificity**

This CNP antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 92-126 amino acids from human CNP.

**Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CNP Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CNP Antibody (N-Term) - Protein Information****Name** CNP ([HGNC:2158](#))

**Function** Catalyzes the formation of 2'-nucleotide products from 2',3'- cyclic substrates (By similarity). May participate in RNA metabolism in the myelinating cell, CNP is the third most abundant protein in central nervous system myelin (By similarity).

### Cellular Location


Membrane {ECO:0000250|UniProtKB:P16330}; Lipid- anchor {ECO:0000250|UniProtKB:P16330}. Melanosome. Note=Firmly bound to membrane structures of brain white matter. {ECO:0000250|UniProtKB:P16330}

### CNP Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CNP Antibody (N-Term) - Images



All lanes : Anti-CNP Antibody (N-Term) at 1:2000 dilution Lane 1: mouse brain lysates Lane 2: rat brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 48 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### CNP Antibody (N-Term) - Background

May participate in RNA metabolism in the myelinating cell, CNP is the third most abundant protein in central nervous system myelin.

### CNP Antibody (N-Term) - References

Thompson R.J.,et al.Biochem. Soc. Trans. 20:621-626(1992).  
Kurihara T.,et al.Biochem. Biophys. Res. Commun. 152:837-842(1988).  
Monoh K.,et al.Gene 129:297-301(1993).  
Douglas A.J.,et al.Ann. Hum. Genet. 56:243-254(1992).  
Zody M.C.,et al.Nature 440:1045-1049(2006).

