

**CYP2E1 Antibody (N-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP21772a****Specification**

---

**CYP2E1 Antibody (N-Term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P05181</a>
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	56849

**CYP2E1 Antibody (N-Term) - Additional Information****Gene ID** 1571**Other Names**

Cytochrome P450 2E1, 11413-, 4-nitrophenol 2-hydroxylase, 11413n7, CYP11E1, Cytochrome P450-J, Cytochrome P450 2E1, N-terminally processed, CYP2E1, CYP2E

**Target/Specificity**

This CYP2E1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 66-100 amino acids from human CYP2E1.

**Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CYP2E1 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CYP2E1 Antibody (N-Term) - Protein Information****Name** CYP2E1 {ECO:0000303|PubMed:10553002, ECO:0000312|HGNC:HGNC:2631}**Function** A cytochrome P450 monooxygenase involved in the metabolism of fatty acids (PubMed:[10553002](#), PubMed:[18577768](#)). Mechanistically, uses molecular oxygen inserting one

oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:[10553002](#), PubMed:[18577768](#)). Catalyzes the hydroxylation of carbon-hydrogen bonds. Hydroxylates fatty acids specifically at the omega-1 position displaying the highest catalytic activity for saturated fatty acids (PubMed:[10553002](#), PubMed:[18577768](#)). May be involved in the oxidative metabolism of xenobiotics (Probable).

#### Cellular Location


Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Microsome membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Note=Post-translationally targeted to mitochondria. TOMM70 is required for the translocation across the mitochondrial outer membrane. After translocation into the matrix, associates with the inner membrane as a membrane extrinsic protein {ECO:0000250|UniProtKB:P05182}

#### CYP2E1 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### CYP2E1 Antibody (N-Term) - Images



Anti-CYP2E1 Antibody (N-Term) at 1:2000 dilution + human liver lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 57 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

#### CYP2E1 Antibody (N-Term) - Background

Metabolizes several precarcinogens, drugs, and solvents to reactive metabolites. Inactivates a number of drugs and xenobiotics and also bioactivates many xenobiotic substrates to their hepatotoxic or carcinogenic forms.

#### **CYP2E1 Antibody (N-Term) - References**

Song B.-J.,et al.J. Biol. Chem. 261:16689-16697(1986).

Umeno M.,et al.Biochemistry 27:9006-9013(1988).

Zhuge J.,et al.Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.

Deloukas P.,et al.Nature 429:375-381(2004).

Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.