

FLNB Antibody (N-Term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP21863a**Specification**

FLNB Antibody (N-Term) - Product Information

Application	WB,E
Primary Accession	O75369
Other Accession	Q80X90
Reactivity	Human, Mouse
Predicted	Mouse
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit IgG
Calculated MW	278164

FLNB Antibody (N-Term) - Additional Information**Gene ID** 2317**Other Names**

Filamin-B, FLN-B, ABP-278, ABP-280 homolog, Actin-binding-like protein, Beta-filamin, Filamin homolog 1, Fh1, Filamin-3, Thyroid autoantigen, Truncated actin-binding protein, Truncated ABP, FLNB, FLN1L, FLN3, TAP, TAP

Target/Specificity

This FLNB antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 288-322 amino acids from human FLNB.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FLNB Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

FLNB Antibody (N-Term) - Protein Information**Name** FLNB

Synonyms FLN1L, FLN3, TABP, TAP

Function Connects cell membrane constituents to the actin cytoskeleton. May promote orthogonal branching of actin filaments and links actin filaments to membrane glycoproteins. Anchors various transmembrane proteins to the actin cytoskeleton. Interaction with FLNA may allow neuroblast migration from the ventricular zone into the cortical plate. Various interactions and localizations of isoforms affect myotube morphology and myogenesis. Isoform 6 accelerates muscle differentiation in vitro.

Cellular Location

[Isoform 1]: Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, stress fiber. Cytoplasm, myofibril, sarcomere, Z line. Note=In differentiating myotubes, isoform 1, isoform 2 and isoform 3 are localized diffusely throughout the cytoplasm with regions of enrichment at the longitudinal actin stress fiber. In differentiated tubes, isoform 1 is also detected within the Z-lines
[Isoform 3]: Cytoplasm, cytoskeleton, stress fiber

Tissue Location


Ubiquitous. Isoform 1 and isoform 2 are expressed in placenta, bone marrow, brain, umbilical vein endothelial cells (HUVEC), retina and skeletal muscle. Isoform 1 is predominantly expressed in prostate, uterus, liver, thyroid, stomach, lymph node, small intestine, spleen, skeletal muscle, kidney, placenta, pancreas, heart, lung, platelets, endothelial cells, megakaryocytic and erythroleukemic cell lines. Isoform 2 is predominantly expressed in spinal cord, platelet and Daudi cells. Also expressed in thyroid adenoma, neurofibrillary tangles (NFT), senile plaques in the hippocampus and cerebral cortex in Alzheimer disease (AD). Isoform 3 and isoform 6 are expressed predominantly in lung, heart, skeletal muscle, testis, spleen, thymus and leukocytes. Isoform 4 and isoform 5 are expressed in heart.

FLNB Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FLNB Antibody (N-Term) - Images



All lanes : Anti-FLNB Antibody (N-Term) at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: A431 whole cell lysate Lane 4: 293T/17 whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 278 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

FLNB Antibody (N-Term) - Background

Connects cell membrane constituents to the actin cytoskeleton. May promote orthogonal branching of actin filaments and links actin filaments to membrane glycoproteins. Anchors various transmembrane proteins to the actin cytoskeleton. Interaction with FLNA may allow neuroblast migration from the ventricular zone into the cortical plate. Various interactions and localizations of isoforms affect myotube morphology and myogenesis. Isoform 6 accelerates muscle differentiation in vitro.

FLNB Antibody (N-Term) - References

Takafuta T.,et al.J. Biol. Chem. 273:17531-17538(1998).
Xu W.-F.,et al.Blood 92:1268-1276(1998).
van Der Flier A.,et al.J. Cell Biol. 156:361-376(2002).
Chakarova C.,et al.Hum. Genet. 107:597-611(2000).
Oshikawa M.,et al.DNA Res. 15:123-136(2008).