

AKR7A3 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP21988c

Specification

AKR7A3 Antibody (Center) - Product Information

Application WB,E
Primary Accession O95154
Reactivity Human
Host Rabbit
Clonality polyclonal
Isotype Rabbit IgG
Calculated MW 37206

AKR7A3 Antibody (Center) - Additional Information

Gene ID 22977

Other Names

Aflatoxin B1 aldehyde reductase member 3, 1.-.-., AFB1 aldehyde reductase 2, AFB1-AR 2, AKR7A3, AFAR2

Target/Specificity

This AKR7A3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 187-217 amino acids from the Central region of human AKR7A3.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

AKR7A3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

AKR7A3 Antibody (Center) - Protein Information

Name AKR7A3

Synonyms AFAR2



Function Can reduce the dialdehyde protein-binding form of aflatoxin B1 (AFB1) to the non-binding AFB1 dialcohol. May be involved in protection of liver against the toxic and carcinogenic effects of AFB1, a potent hepatocarcinogen.

Cellular Location Cytoplasm.

Tissue Location


Expressed in colon, kidney, liver, pancreas, adenocarcinoma and endometrium.

AKR7A3 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

AKR7A3 Antibody (Center) - Images




All lanes : Anti-AKR7A3 Antibody (Center) at 1:2000 dilution Lane 1: human liver lysate Lane 2: SK-BR-3 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

AKR7A3 Antibody (Center) - Background

Can reduce the dialdehyde protein-binding form of aflatoxin B1 (AFB1) to the non-binding AFB1 dialcohol. May be involved in protection of liver against the toxic and carcinogenic effects of AFB1, a potent hepatocarcinogen.

AKR7A3 Antibody (Center) - References





Knight L.P., et al. Carcinogenesis 20:1215-1223(1999).

Praml C., et al. Oncogene 22:4765-4773(2003).

Gregory S.G., et al. Nature 441:315-321(2006).

Bodreddigari S., et al. Chem. Res. Toxicol. 21:1134-1142(2008).