

**IL4I1 Antibody (C-Term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP22006b****Specification**

---

**IL4I1 Antibody (C-Term) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB,E                   |
| Primary Accession | <a href="#">O96RQ9</a> |
| Reactivity        | Human                  |
| Host              | Rabbit                 |
| Clonality         | polyclonal             |
| Isotype           | Rabbit IgG             |

**IL4I1 Antibody (C-Term) - Additional Information****Gene ID** 259307**Other Names**

L-amino-acid oxidase, LAAO, LAO, 1.4.3.2, Interleukin-4-induced protein 1, IL4-induced protein 1, Protein Fig-1, hFIG1, IL4I1, FIG1

**Target/Specificity**

This IL4I1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 391-422 amino acids from human IL4I1.

**Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

IL4I1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**IL4I1 Antibody (C-Term) - Protein Information****Name** IL4I1 {ECO:0000303|PubMed:16029492}

**Function** Secreted L-amino-acid oxidase that acts as a key immunoregulator (PubMed:[17356132](#), PubMed:[32818467](#), PubMed:[32866000](#)). Has preference for L-aromatic amino acids: converts phenylalanine (Phe), tyrosine (Tyr) and tryptophan (Trp) to phenylpyruvic acid (PP),

hydroxyphenylpyruvic acid (HPP), and indole-3-pyruvic acid (I3P), respectively (PubMed:[17356132](#), PubMed:[32818467](#), PubMed:[32866000](#)). Also has weak L-arginine oxidase activity (PubMed:[26673964](#)). Acts as a negative regulator of anti-tumor immunity by mediating Trp degradation via an indole pyruvate pathway that activates the transcription factor AHR (PubMed:[32818467](#), PubMed:[32866000](#)). IL4I1-mediated Trp catabolism generates I3P, giving rise to indole metabolites (indole-3-acetic acid (IAA) and indole-3-aldehyde (I3A)) and kynurenic acid, which act as ligands for AHR, a ligand-activated transcription factor that plays important roles in immunity and cancer (PubMed:[32818467](#), PubMed:[32866000](#)). AHR activation by indoles following IL4I1-mediated Trp degradation enhances tumor progression by promoting cancer cell motility and suppressing adaptive immunity (PubMed:[32818467](#)). Also has an immunoregulatory function in some immune cells, probably by mediating Trp degradation and promoting downstream AHR activation: inhibits T-cell activation and proliferation, promotes the differentiation of naive CD4(+) T-cells into FOXP3(+) regulatory T- cells (Treg) and regulates the development and function of B-cells (PubMed:[17356132](#), PubMed:[25446972](#), PubMed:[25778793](#), PubMed:[28891065](#)). Also regulates M2 macrophage polarization by inhibiting T-cell activation (By similarity). Also has antibacterial properties by inhibiting growth of Gram negative and Gram positive bacteria through the production of NH<sub>4</sub>(+) and H<sub>2</sub>O<sub>2</sub> (PubMed:[23355881](#)).

#### **Cellular Location**

Secreted. Lysosome {ECO:0000250|UniProtKB:O09046}. Cytoplasmic vesicle, secretory vesicle, acrosome. Note=Secreted at the immunological synapse.

#### **Tissue Location**

Primarily found in immune tissues, with the highest expression in lymph nodes and spleen (PubMed:12031486, PubMed:12446450). Present in germinal center macrophages and inflammatory myeloid cells and antigen-presenting cells (at protein level) (PubMed:17356132). Also present in spermatozoa (at protein level) (PubMed:25767141). Highly expressed in primary mediastinal large B-cell lymphoma, a specific subtype of diffuse large B-cell lymphoma (PubMed:12446450). Expressed by neoplastic cells of several B-cell lymphomas and by tumor-associated macrophages (PubMed:19436310)


#### **IL4I1 Antibody (C-Term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **IL4I1 Antibody (C-Term) - Images**





All lanes : Anti-IL4I1 Antibody (C-Term) at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: HT-1080 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 63 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

#### **IL4I1 Antibody (C-Term) - Background**

Lysosomal L-amino-acid oxidase with highest specific activity with phenylalanine. May play a role in lysosomal antigen processing and presentation (By similarity).

#### **IL4I1 Antibody (C-Term) - References**

Chavan S.S.,et al.Biochim. Biophys. Acta 1576:70-80(2002).  
Wiemann S.,et al.BMC Biol. 3:16-16(2005).  
Chu C.C.,et al.Submitted (MAY-2005) to the EMBL/GenBank/DDBJ databases.  
Jikuya H.,et al.Submitted (JAN-2002) to the EMBL/GenBank/DDBJ databases.  
Clark H.F.,et al.Genome Res. 13:2265-2270(2003).

#### **IL4I1 Antibody (C-Term) - Citations**

- [High expression of IL4I1 is correlated with poor prognosis and immune infiltration in thyroid cancer](#)