

RPS7 Antibody (N-Term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP22077a

Specification

RPS7 Antibody (N-Term) - Product Information

Application Primary Accession Other Accession

Reactivity Predicted Host Clonality Isotype Calculated MW WB, FC,E <u>P62081</u> A6H769, <u>P62084</u>, <u>05RT64</u>, <u>P62082</u>, <u>P62083</u>, <u>P50894</u>, <u>P02362</u>, <u>G1SVB0</u> Human, Mouse Bovine, Zebrafish, Rabbit, Rat, Xenopus Rabbit polyclonal Rabbit IgG 22127

RPS7 Antibody (N-Term) - Additional Information

Gene ID 6201

Other Names 40S ribosomal protein S7, RPS7

Target/Specificity

This RPS7 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 7-40 amino acids from human RPS7.

Dilution WB~~1:2000 FC~~1:25 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions RPS7 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

RPS7 Antibody (N-Term) - Protein Information

Name RPS7 (<u>HGNC:10440</u>)



Function Component of the small ribosomal subunit (PubMed:<u>23636399</u>). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed:<u>23636399</u>). Required for rRNA maturation (PubMed:<u>19061985</u>). Part of the small subunit (SSU) processome, first precursor of the small eukaryotic ribosomal subunit. During the assembly of the SSU processome in the nucleolus, many ribosome biogenesis factors, an RNA chaperone and ribosomal proteins associate with the nascent pre-rRNA and work in concert to generate RNA folding, modifications, rearrangements and cleavage as well as targeted degradation of pre-ribosomal RNA by the RNA exosome (PubMed:<u>34516797</u>).

Cellular Location


Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm. Nucleus, nucleolus Note=Although RPS7 is functional within the cytoplasm, the assembly of ribosomal subunits occurs in the nucleus. RPS7 nuclear import is mediated by IPO5/RanBP5, IPO7/RanBP7, KPNB1/importin-beta or TPNO1/Trn (PubMed:9687515). Colocalizes with NEK6 in the centrosome (PubMed:20873783).

RPS7 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- <u>Cell Culture</u>

RPS7 Antibody (N-Term) - Images



All lanes : Anti-RPS7 Antibody (N-Term) at 1:2000 dilution Lane 1: Hela whole cell lysate Lane 2: A549 whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: Jurkat whole cell lysate Lane 5: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit lgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 22 kDa Blocking/Dilution buffer: 5% NFDM/TBST.





Overlay histogram showing Hela cells stained with AP22077a (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then icubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP22077a, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit lgG, **DyLight**® 488 Conjugated Highly Cross-Adsorbed(OH191631) at 1/200 dilution for 40 min at 37ºC. Isotype control antibody (blue line) was rabbit IgG $(1\mu g/1 \times 10^{6} \text{ cells})$ used under the same conditions. Acquisition of >10, 000 events was performed.

RPS7 Antibody (N-Term) - Background

Required for rRNA maturation.

RPS7 Antibody (N-Term) - References

Annilo T., et al.Gene 165:297-302(1995). Ota T., et al.Nat. Genet. 36:40-45(2004). Hillier L.W., et al.Nature 434:724-731(2005). Mural R.J., et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Vladimirov S.N., et al.Eur. J. Biochem. 239:144-149(1996).