

CD56 antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP22454b**Specification**

CD56 antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P13591
Reactivity	Human
Host	Rabbit
Clonality	polyclonal
Isotype	Rabbit Ig
Calculated MW	94574

CD56 antibody (C-term) - Additional Information**Gene ID** 4684**Other Names**

Neural cell adhesion molecule 1, N-CAM-1, NCAM-1, CD56, NCAM1 (HGNC:7656), NCAM

Target/Specificity

This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between amino acids from human.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CD56 antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CD56 antibody (C-term) - Protein Information**Name** NCAM1 ([HGNC:7656](#))**Synonyms** NCAM

Function This protein is a cell adhesion molecule involved in neuron- neuron adhesion, neurite fasciculation, outgrowth of neurites, etc. (Microbial infection) Acts as a receptor for Zika virus.

Cellular Location


[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Isoform 3]: Cell membrane; Lipid-anchor, GPI- anchor [Isoform 5]: Secreted.

CD56 antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD56 antibody (C-term) - Images



All lanes: Anti-CD56 antibody (C-term) at 1:1000 dilution Lane 1: U-87 MG whole cell lysate Lane 2: SW480 whole cell lysate Lane 3: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 150 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

CD56 antibody (C-term) - Background

This protein is a cell adhesion molecule involved in neuron- neuron adhesion, neurite fasciculation, outgrowth of neurites, etc. (Microbial infection) Acts as a receptor for Zika virus.

CD56 antibody (C-term) - References

Barton C.H.,et al.Development 104:165-173(1988).
Lanier L.L.,et al.J. Immunol. 146:4421-4426(1991).
Saito S.,et al.Lung Cancer 10:307-318(1994).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Totoki Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.