

MARCKS Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2521b**Specification**

MARCKS Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P29966
Other Accession	NP_002347
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	31555
Antigen Region	241-271

MARCKS Antibody (C-term) - Additional Information**Gene ID** 4082**Other Names**

Myristoylated alanine-rich C-kinase substrate, MARCKS, Protein kinase C substrate, 80 kDa protein, light chain, 80K-L protein, PKCSL, MARCKS, MACS, PRKCSL

Target/Specificity

This MARCKS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 241-271 amino acids from the C-terminal region of human MARCKS.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MARCKS Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MARCKS Antibody (C-term) - Protein Information**Name** MARCKS

Synonyms MACS, PRKCSL

Function Membrane-associated protein that plays a role in the structural modulation of the actin cytoskeleton, chemotaxis, motility, cell adhesion, phagocytosis, and exocytosis through lipid sequestering and/or protein docking to membranes (PubMed:[23704996](#), PubMed:[36009319](#)). Thus, exerts an influence on a plethora of physiological processes, such as embryonic development, tissue regeneration, neuronal plasticity, and inflammation. Sequesters phosphatidylinositol 4,5-bisphosphate (PIP2) at lipid rafts in the plasma membrane of quiescent cells, an action reversed by protein kinase C, ultimately inhibiting exocytosis (PubMed:[23704996](#)). During inflammation, promotes the migration and adhesion of inflammatory cells and the secretion of cytokines such as tumor necrosis factor (TNF), particularly in macrophages (PubMed:[37949888](#)). Plays an essential role in bacteria- induced intracellular reactive oxygen species (ROS) formation in the monocytic cell type. Participates in the regulation of neurite initiation and outgrowth by interacting with components of cellular machinery including CDC42 that regulates cell shape and process extension through modulation of the cytoskeleton (By similarity). Plays also a role in axon development by mediating docking and fusion of RAB10-positive vesicles with the plasma membrane (By similarity).

Cellular Location

Cell membrane; Lipid-anchor. Cytoplasm, cytoskeleton Cytoplasm. Note=PKC-dependent phosphorylation displaces MARCKS from the cell membrane and subsequent dephosphorylation is accompanied by its reassociation with the membrane.


Tissue Location

Detected in spermatozoa.

MARCKS Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MARCKS Antibody (C-term) - Images

Western blot analysis of hMARCKS-K254 (Cat. #AP2521b) in Hela cell line lysates (35ug/lane). MARCKS (arrow) was detected using the purified Pab.

MARCKS Antibody (C-term) - Background

MARCKS is a substrate for protein kinase C. It is localized to the plasma membrane and is an actin filament crosslinking protein. Phosphorylation by protein kinase C or binding to calcium-calmodulin inhibits its association with actin and with the plasma membrane, leading to its presence in the cytoplasm. The protein is thought to be involved in cell motility, phagocytosis, membrane trafficking and mitogenesis.

MARCKS Antibody (C-term) - References

Rauch, M.E., et al., J. Biol. Chem. 277(16):14068-14076 (2002).
Aderem, A., Biochem. Soc. Trans. 23(3):587-591 (1995).
Rao, P.H., et al., Cytogenet. Cell Genet. 66(4):272-273 (1994).
Taniguchi, H., et al., J. Biol. Chem. 268(14):9960-9963 (1993).
Blackshear, P.J., J. Biol. Chem. 268(3):1501-1504 (1993).