

## **CPT1B Antibody (C-term)**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2532b

## **Specification**

# CPT1B Antibody (C-term) - Product Information

**Application** WB, IHC-P,E **Primary Accession** 092523 NP 004368 Other Accession Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 87801 Antigen Region 606-637

## CPT1B Antibody (C-term) - Additional Information

#### **Gene ID 1375**

#### **Other Names**

Carnitine O-palmitoyltransferase 1, muscle isoform, CPT1-M, Carnitine O-palmitoyltransferase I, muscle isoform, CPT I, CPTI-M, Carnitine palmitoyltransferase 1B, Carnitine palmitoyltransferase I-like protein, CPT1B, KIAA1670

## Target/Specificity

This CPT1B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 606-637 amino acids from the C-terminal region of human CPT1B.

#### **Dilution**

WB~~1:1000 IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

CPT1B Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### **CPT1B Antibody (C-term) - Protein Information**



## Name CPT1B

# Synonyms KIAA1670

**Function** Catalyzes the transfer of the acyl group of long-chain fatty acid-CoA conjugates onto carnitine, an essential step for the mitochondrial uptake of long-chain fatty acids and their subsequent beta-oxidation in the mitochondrion.

## **Cellular Location**

Mitochondrion outer membrane; Multi-pass membrane protein

#### **Tissue Location**


Strong expression in heart and skeletal muscle. No expression in liver and kidney

## **CPT1B Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# CPT1B Antibody (C-term) - Images



Western blot analysis of CPT1B (arrow) using rabbit polyclonal CPT1B Antibody (C-term) (Cat.#AP2532b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the CPT1B gene (Lane 2).





Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

# CPT1B Antibody (C-term) - Background

CPT1B is a member of the carnitine/choline acetyltransferase family, and is the rate-controlling enzyme of the long-chain fatty acid beta-oxidation pathway in muscle mitochondria. This enzyme is required for the net transport of long-chain fatty acyl-CoAs from the cytoplasm into the mitochondria.

## **CPT1B Antibody (C-term) - References**

Dai, J., et al., Biochem. Biophys. Res. Commun. 301(3):758-763 (2003). van der Leij, F.R., et al., J. Biol. Chem. 277(30):26994-27005 (2002). Hirosawa, M., et al., DNA Res. 8(1):1-9 (2001). Yamazaki, N., et al., J. Biol. Chem. 275(41):31739-31746 (2000). Britton, C.H., et al., Genomics 40(1):209-211 (1997).