

B2M Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2771A**Specification**

B2M Antibody (N-term) - Product Information

Application	IHC-P, FC, WB,E
Primary Accession	P61769
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	13715
Antigen Region	8-39

B2M Antibody (N-term) - Additional Information**Gene ID** 567**Other Names**

Beta-2-microglobulin, Beta-2-microglobulin form pl 53, B2M

Target/Specificity

This B2M antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 8-39 amino acids from the N-terminal region of human B2M.

Dilution

IHC-P~~1:10~50

FC~~1:10~50

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

B2M Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

B2M Antibody (N-term) - Protein Information**Name** B2M ([HGNC:914](#))

Function Component of the class I major histocompatibility complex (MHC). Involved in the presentation of peptide antigens to the immune system. Exogenously applied M.tuberculosis EsxA or EsxA-EsxB (or EsxA expressed in host) binds B2M and decreases its export to the cell surface (total protein levels do not change), probably leading to defects in class I antigen presentation (PubMed:[25356553](#)).

Cellular Location


Secreted. Cell surface. Note=Detected in serum and urine (PubMed:1336137, PubMed:7554280). {ECO:0000269|PubMed:7554280, ECO:0000269|Ref.6}

B2M Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


B2M Antibody (N-term) - Images



Anti-B2M Antibody (N-term) at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 14 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with B2M antibody (N-term) (Cat.#AP2771a), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



B2M Antibody (N-term) (Cat. #AP2771a) flow cytometric analysis of Hela cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

B2M Antibody (N-term) - Background

Beta-2-microglobulin is a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells.

B2M Antibody (N-term) - References

- Sakata, M., J. Mol. Biol. 382 (5), 1242-1255 (2008)
- Huang, W.C., Clin. Cancer Res. 14 (17), 5341-5347 (2008)
- Gattoni-Celli, S., Cancer Res. 52 (5), 1201-1204 (1992)