

PRDX1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP2924c

Specification

PRDX1 Antibody (Center) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region IHC-P, FC, WB,E <u>Q06830</u> <u>Q63716</u>, <u>Q9JKY1</u> Human Hamster, Rat Rabbit Polyclonal Rabbit IgG 22110 88-116

PRDX1 Antibody (Center) - Additional Information

Gene ID 5052

Other Names

Peroxiredoxin-1, Natural killer cell-enhancing factor A, NKEF-A, Proliferation-associated gene protein, PAG, Thioredoxin peroxidase 2, Thioredoxin-dependent peroxide reductase 2, PRDX1, PAGA, PAGB, TDPX2

Target/Specificity

This PRDX1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 88-116 amino acids from the Central region of human PRDX1.

Dilution IHC-P~~1:50~100 FC~~1:10~50 WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PRDX1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

PRDX1 Antibody (Center) - Protein Information



Name PRDX1

Synonyms PAGA, PAGB, TDPX2

Function Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H(2)O(2) (PubMed:<u>9497357</u>). Reduces an intramolecular disulfide bond in GDPD5 that gates the ability to GDPD5 to drive postmitotic motor neuron differentiation (By similarity).

Cellular Location

Cytoplasm. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

PRDX1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

PRDX1 Antibody (Center) - Images


-
1
•

Western blot analysis of PRDX1 Antibody (Center) (Cat. #AP2924c) in CEM cell line lysates (35ug/lane). PRDX1 (arrow) was detected using the purified Pab.






Western blot analysis of PRDX1 (arrow) using rabbit polyclonal PRDX1 Antibody (Center) (Cat. #AP2924c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the PRDX1 gene (Lane 2).



PRDX1 Antibody (Center) (Cat. #AP2924c) immunohistochemistry analysis in formalin fixed and paraffin embedded human colon carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the PRDX1 Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



PRDX1 Antibody (Center) (Cat. #AP2924c) flow cytometric analysis of 293 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PRDX1 Antibody (Center) - Background

PRDX1 is a member of the peroxiredoxin family of antioxidant enzymes, which reduce hydrogen



peroxide and alkyl hydroperoxides. This protein may play an antioxidant protective role in cells, and may contribute to the antiviral activity of CD8(+) T-cells. This protein may have a proliferative effect and play a role in cancer development or progression.

PRDX1 Antibody (Center) - References

Song,I.S., et.al., Carcinogenesis 30 (7), 1106-1114 (2009)