

PRDX3 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP2926a

Specification

PRDX3 Antibody (N-term) - Product Information

Application	WB, FC, IHC-P,E
Primary Accession	P30048
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	27693
Antigen Region	66-94

PRDX3 Antibody (N-term) - Additional Information

Gene ID 10935

Other Names

Thioredoxin-dependent peroxide reductase, mitochondrial, Antioxidant protein 1, AOP-1, HBC189, Peroxiredoxin III, Prx-III, Peroxiredoxin-3, Protein MER5 homolog, PRDX3, AOP1

Target/Specificity

This PRDX3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 66-94 amino acids from the N-terminal region of human PRDX3.

Dilution

WB~~1:1000

FC~~1:10~50

IHC-P~~1:50~100

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PRDX3 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

PRDX3 Antibody (N-term) - Protein Information

Name PRDX3

Synonyms AOP1

Function Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative stress by detoxifying peroxides (PubMed:[17707404](#), PubMed:[29438714](#), PubMed:[33889951](#), PubMed:[7733872](#)). Acts synergistically with MAP3K13 to regulate the activation of NF-kappa-B in the cytosol (PubMed:[12492477](#)). Required for the maintenance of physical strength (By similarity).

Cellular Location


Mitochondrion. Cytoplasm. Early endosome. Note=Localizes to early endosomes in a RPS6KC1-dependent manner.

PRDX3 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PRDX3 Antibody (N-term) - Images




Western blot analysis of PRDX3 Antibody (N-term) (Cat. #AP2926a) in NCI-H460,MDA-MB231,HepG2 cell line lysates (35ug/lane). PRDX3 (arrow) was detected using the purified Pab.(2ug/ml)



Western blot analysis of PRDX3 (arrow) using rabbit polyclonal PRDX3 Antibody (N-term) (Cat. #AP2926a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the PRDX3 gene.



Formalin-fixed and paraffin-embedded human lung carcinoma reacted with PRDX3 Antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of MDA-MB468 cells using PRDX3 Antibody (N-term)(bottom histogram) compared to a negative control cell (top histogram) FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

PRDX3 Antibody (N-term) - Background

PRDX3 is a protein with antioxidant function and is localized in the mitochondrion. This protein is

involved in redox regulation of the cell and Protects radical-sensitive enzymes from oxidative damage by a radical-generating system. It Acts synergistically with MAP3K13 to regulate the activation of NF-kappa-B in the cytosol.

PRDX3 Antibody (N-term) - References

Cox, A.G., et.al., Biochemistry 48 (27), 6495-6501 (2009)