

## Phospho-PDHE1A(S232) Antibody

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP3758A

## **Specification**

## Phospho-PDHE1A(S232) Antibody - Product Information

Application WB, DB,E
Primary Accession P08559

Other Accession <u>P26284</u>, <u>P29804</u>, <u>P35486</u>, <u>Q8HXW9</u>, <u>A7MB35</u>,

NP\_000275.1

Reactivity Human

Predicted Bovine, Monkey, Mouse, Pig, Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG

## Phospho-PDHE1A(S232) Antibody - Additional Information

#### **Gene ID 5160**

#### **Other Names**

Pyruvate dehydrogenase E1 component subunit alpha, somatic form, mitochondrial, PDHE1-A type I, PDHA1, PHE1A

## Target/Specificity

This PDHE1A Antibody is generated from rabbits immunized with a KLH conjugated synthetic phosphopeptide corresponding to amino acid residues surrounding S232 of human PDHE1A.

## **Dilution**

WB~~1:5000 DB~~1:500

E~~Use at an assay dependent concentration.

## **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

Phospho-PDHE1A(S232) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Phospho-PDHE1A(S232) Antibody - Protein Information

## Name PDHA1





Tel: 858.875.1900 Fax: 858.875.1999

## Synonyms PHE1A

Function The pyruvate dehydrogenase complex catalyzes the overall conversion of pyruvate to acetyl-CoA and CO(2), and thereby links the glycolytic pathway to the tricarboxylic cycle.

**Cellular Location** Mitochondrion matrix.


**Tissue Location** Ubiquitous.

## Phospho-PDHE1A(S232) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# Phospho-PDHE1A(S232) Antibody - Images




Western blot analysis of lysates from C6 cell line, untreated or treated with CIP, 100ng/ml, using Phospho-PDHE1A(S232) Antibody(upper) or Tubulin (lower).





Western blot analysis of lysates from 293 cell line, untreated or treated with CIP, 100ng/ml, using Phospho-PDHE1A(S232) Antibody(upper) or Tubulin (lower).



Dot blot analysis of anti-mPDHE1A-S232 Phospho-specific Pab (Cat. #AP3758a) on nitrocellulose membrane. 50ng of Phospho-peptide or Non Phospho-peptide per dot were adsorbed. Antibody working concentrations are 0.6ug per ml.

## Phospho-PDHE1A(S232) Antibody - Citations

- Energy sensor AMPK gamma regulates translation via phosphatase PPP6C independent of AMPK alpha
- PINK1 Is a Negative Regulator of Growth and the Warburg Effect in Glioblastoma.
- JX06 Selectively Inhibits Pyruvate Dehydrogenase Kinase PDK1 by a Covalent Cysteine Modification.