

### **RAB27B Antibody (C-term)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP4986b

## **Specification**

# RAB27B Antibody (C-term) - Product Information

**Application** FC, WB, E **Primary Accession** 000194 NP 004154 Other Accession Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 24608 Antigen Region 186-215

### RAB27B Antibody (C-term) - Additional Information

**Gene ID 5874** 

### **Other Names**

Ras-related protein Rab-27B, C25KG, RAB27B

### Target/Specificity

This RAB27B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 186-215 amino acids from the C-terminal region of human RAB27B.

# **Dilution**

FC~~1:10~50 WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

RAB27B Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# RAB27B Antibody (C-term) - Protein Information

Name RAB27B





Tel: 858.875.1900 Fax: 858.875.1999

Function Small GTPase which cycles between active GTP-bound and inactive GDP-bound states. In its active state, binds to a variety of effector proteins to regulate homeostasis of late endocytic pathway, including endosomal positioning, maturation and secretion (PubMed: 30771381). Plays a role in NTRK2/TRKB axonal anterograde transport by facilitating the association of NTRK2/TRKB with KLC1 (PubMed: 21775604). May be involved in targeting uroplakins to urothelial apical membranes (By similarity).

### **Cellular Location**

Membrane; Lipid-anchor. Late endosome

### **Tissue Location**


Expressed primarily in testis.

### RAB27B Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## RAB27B Antibody (C-term) - Images



Western blot analysis of RAB27B Antibody (C-term) (Cat. #AP4986b) in NCI-H460 cell line lysates (35ug/lane). RAB27B (arrow) was detected using the purified Pab.





RAB27B Antibody (C-term) (Cat. #AP4986b) flow cytometric analysis of NCI-H460 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

# RAB27B Antibody (C-term) - Background

RAB27B may be involved in targeting uroplakins to urothelial apical membranes.

## RAB27B Antibody (C-term) - References

Hayasaka, S., et al. Asian J. Androl. 10(4):561-568(2008) Shirakawa, R., et al. J. Biol. Chem. 279(11):10730-10737(2004) Tiwari, S., et al. Blood 102(12):3970-3979(2003)