

CDK5RAP3 Antibody
Affinity-Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP50663**Specification**

CDK5RAP3 Antibody - Product Information

Application	WB
Primary Accession	O96JB5
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57,46,31 KDa
Antigen Region	350-379

CDK5RAP3 Antibody - Additional Information**Gene ID** 80279**Other Names**

CDK5 regulatory subunit-associated protein 3, CDK5 activator-binding protein C53, Protein HSF-27, CDK5RAP3, IC53

Dilution

WB~~1:500

FormatRabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.**Storage Conditions**

-20°C

CDK5RAP3 Antibody - Protein Information**Name** CDK5RAP3 {ECO:0000303|PubMed:30635284, ECO:0000312|HGNC:HGNC:18673}**Function**

Substrate adapter of E3 ligase complexes mediating ufmylation, the covalent attachment of the ubiquitin-like modifier UFM1 to substrate proteins, and which is involved in various processes, such as ribosome recycling and reticulophagy (also called ER-phagy) (PubMed:23152784, PubMed:30635284, PubMed:32851973, PubMed:36121123, PubMed:36543799, PubMed:37595036, PubMed:38383785, PubMed:38383789). As part of

the UREL complex, plays a key role in ribosome recycling by promoting mono-ufmylation of RPL26/uL24 subunit of the 60S ribosome (PubMed:38383785, PubMed:38383789). Ufmylation of RPL26/uL24 occurs on free 60S ribosomes following ribosome dissociation: it weakens the junction between post-termination 60S subunits and SEC61 translocons, promoting release and recycling of the large ribosomal subunit from the endoplasmic reticulum membrane (PubMed:38383785, PubMed:38383789). Ufmylation of RPL26/uL24 and subsequent 60S ribosome recycling either take place after normal termination of translation or after ribosome stalling during cotranslational translocation at the endoplasmic reticulum (PubMed:32851973, PubMed:37595036, PubMed:38383785, PubMed:38383789). Within the UREL complex, CDK5RAP3 acts as a substrate adapter that constrains UFL1 ligase activity to mono-ufmylate RPL26/uL24 at 'Lys-134' (PubMed:36121123, PubMed:38383785, PubMed:38383789). The UREL complex is also involved in reticulophagy in response to endoplasmic reticulum stress by promoting ufmylation of proteins such as CYB5R3, thereby promoting lysosomal degradation of ufmylated proteins (PubMed:36543799). Also acts as a regulator of transcription: negatively regulates NF-kappa-B-mediated gene transcription through the control of RELA phosphorylation (PubMed:17785205, PubMed:20228063). Also regulates mitotic G2/M transition checkpoint and mitotic G2 DNA damage checkpoint (PubMed:15790566, PubMed:19223857). Through its interaction with CDKN2A/ARF and MDM2 may induce MDM2-dependent p53/TP53 ubiquitination, stabilization and activation in the nucleus, thereby promoting G1 cell cycle arrest and inhibition of cell proliferation (PubMed:16173922). May also play a role in the rupture of the nuclear envelope during apoptosis (PubMed:23478299). May regulate MAPK14 activity by regulating its dephosphorylation by PPM1D/WIP1 (PubMed:21283629). Required for liver development (By similarity).

Cellular Location

Endoplasmic reticulum membrane. Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton. Note=Tethered to the endoplasmic reticulum membrane as part of the UFM1 ribosome E3 ligase (UREL) complex (PubMed:38383785, PubMed:38383789). Colocalizes and associates with microtubules (PubMed:23478299)

Tissue Location

Ubiquitously expressed (PubMed:10721722, PubMed:12054757). Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Isoform 3 is expressed in kidney, liver, skeletal muscle and placenta (PubMed:12737517)


CDK5RAP3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDK5RAP3 Antibody - Images



Western blot analysis of lysate from 293T cell line, using CDK5RAP3 Antibody (AP50663). AP50663 was diluted at 1:500. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

CDK5RAP3 Antibody - Background

Potential regulator of CDK5 activity. May be involved in cell proliferation. Regulates CDK5 activity via its interaction with CDK5R1 (By similarity).

CDK5RAP3 Antibody - References

- Chen J., et al. Biochem. Biophys. Res. Commun. 294:161-166(2002).
Xie Y.H., et al. Cell Res. 13:83-91(2003).
Favier A.-L., et al. Submitted (JAN-2001) to the EMBL/GenBank/DDBJ databases.
Shichijo S., et al. Submitted (MAY-2001) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).