

ADIPOR2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5105C**Specification**

ADIPOR2 Antibody (Center) - Product Information

Application	IF, WB, FC,E
Primary Accession	Q86V24
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	45-72

ADIPOR2 Antibody (Center) - Additional Information**Gene ID** 79602**Other Names**

Adiponectin receptor protein 2, Progestin and adipoQ receptor family member II, ADIPOR2, PAQR2

Target/Specificity

This ADIPOR2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 45-72 amino acids from the Central region of human ADIPOR2.

Dilution

IF~~1:10~50

WB~~1:1000

FC~~1:25

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ADIPOR2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ADIPOR2 Antibody (Center) - Protein Information**Name** ADIPOR2 ([HGNC:24041](#))**Function** Receptor for ADIPOQ, an essential hormone secreted by adipocytes that regulates

glucose and lipid metabolism (PubMed:[12802337](#), PubMed:[25855295](#)). Required for normal body fat and glucose homeostasis. ADIPOQ-binding activates a signaling cascade that leads to increased PPARA activity, and ultimately to increased fatty acid oxidation and glucose uptake. Has intermediate affinity for globular and full-length adiponectin. Required for normal revascularization after chronic ischemia caused by severing of blood vessels (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein Note=Localized to the cell membrane and intracellular organelles

Tissue Location


Ubiquitous (PubMed:16044242). Highly expressed in skeletal muscle, liver and placenta (PubMed:12802337). Weakly expressed in brain, heart, colon, spleen, kidney, thymus, small intestine, peripheral blood leukocytes and lung (PubMed:12802337)

ADIPOR2 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ADIPOR2 Antibody (Center) - Images




Confocal immunofluorescent analysis of ADIPOR2 Antibody (Center) (Cat. #AP5105c) with Hela cell followed by Alexa Fluor[®] 489-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).




Western blot analysis of ADIPOR2 Antibody (Center) (Cat. #AP5105c) in Hela cell line lysates (35ug/lane).ADIPOR2 (arrow) was detected using the purified Pab.




All lanes : Anti-ADIPOR2 Antibody (Center) at 1:2000 dilution Lane 1: U-251 MG whole cell lysate Lane 2: U-2OS whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



Anti-ADIPOR2 Antibody (Center) at 1:1000 dilution + Human liver tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-ADIPOR2 Antibody (Center) at 1:1000 dilution + Human skeletal muscle tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing U-2 OS cells stained with AP5105c(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP5105c, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OE188374) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

ADIPOR2 Antibody (Center) - Background

The adiponectin receptors, ADIPOR1 (MIM 607945) and ADIPOR2, serve as receptors for globular and full-length adiponectin (MIM 605441) and mediate increased AMPK (see MIM 602739) and PPAR-alpha (PPARA; MIM 170998) ligand activities, as well as fatty acid oxidation and glucose uptake by adiponectin.

ADIPOR2 Antibody (Center) - References

Ferguson, J.F., et al. Am. J. Clin. Nutr. 91(3):794-801(2010) Halvatsiotis, I., et al. Cardiovasc Diabetol 9, 10 (2010) Yoshihara, K., et al. J Reprod Med 54 (11-12), 669-674 (2009)

ADIPOR2 Antibody (Center) - Citations

- [Contrasting Effects of Adipokines on the Cytokine Production by Primary Human Bronchial Epithelial Cells: Inhibitory Effects of Adiponectin](#)
- [Novel phytopeptide osmotin mimics preventive effects of adiponectin on vascular inflammation and atherosclerosis.](#)