

SHP-1 (Phospho-Tyr564) Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP52495**Specification**

SHP-1 (Phospho-Tyr564) Antibody - Product Information

Application	WB
Primary Accession	P29350
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67561

SHP-1 (Phospho-Tyr564) Antibody - Additional Information**Gene ID** 5777**Other Names**

Tyrosine-protein phosphatase non-receptor type 6, Hematopoietic cell protein-tyrosine phosphatase, Protein-tyrosine phosphatase 1C, PTP-1C, Protein-tyrosine phosphatase SHP-1, SH-PTP1, PTPN6, HCP, PTP1C

Dilution

WB~~1:1000

Format

Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions

-20°C

SHP-1 (Phospho-Tyr564) Antibody - Protein Information**Name** PTPN6**Synonyms** HCP, PTP1C**Function**

Tyrosine phosphatase enzyme that plays important roles in controlling immune signaling pathways and fundamental physiological processes such as hematopoiesis (PubMed:14739280, PubMed:29925997). Dephosphorylates and negatively regulate several receptor tyrosine kinases (RTKs) such as EGFR, PDGFR and FGFR, thereby modulating their signaling activities (PubMed:21258366, PubMed:9733788). When recruited to immunoreceptor tyrosine-based inhibitory motif (ITIM)-containing receptors such as

immunoglobulin-like transcript 2/LILRB1, programmed cell death protein 1/PDCD1, CD3D, CD22, CLEC12A and other receptors involved in immune regulation, initiates their dephosphorylation and subsequently inhibits downstream signaling events (PubMed:11907092, PubMed:14739280, PubMed:37932456, PubMed:38166031). Modulates the signaling of several cytokine receptors including IL-4 receptor (PubMed:9065461). Additionally, targets multiple cytoplasmic signaling molecules including STING1, LCK or STAT1 among others involved in diverse cellular processes including modulation of T-cell activation or cGAS-STING signaling (PubMed:34811497, PubMed:38532423). Within the nucleus, negatively regulates the activity of some transcription factors such as NFAT5 via direct dephosphorylation. Also acts as a key transcriptional regulator of hepatic gluconeogenesis by controlling recruitment of RNA polymerase II to the PCK1 promoter together with STAT5A (PubMed:37595871).

Cellular Location

Cytoplasm. Nucleus Note=In neurons, translocates into the nucleus after treatment with angiotensin II (By similarity). Shuttles between the cytoplasm and nucleus via its association with PDPK1.

Tissue Location


Isoform 1 is expressed in hematopoietic cells. Isoform 2 is expressed in non-hematopoietic cells


SHP-1 (Phospho-Tyr564) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SHP-1 (Phospho-Tyr564) Antibody - Images





Western blot analysis of extracts from COS7 cells, treated with EGF (200ng/ml, 30mins), using SHP-1 (Phospho-Tyr564) antibody.

SHP-1 (Phospho-Tyr564) Antibody - Background

Modulates signaling by tyrosine phosphorylated cell surface receptors such as KIT and the EGF receptor/EGFR. The SH2 regions may interact with other cellular components to modulate its own phosphatase activity against interacting substrates. Together with MTUS1, induces UBE2V2 expression upon angiotensin II stimulation. Plays a key role in hematopoiesis.

SHP-1 (Phospho-Tyr564) Antibody - References

Yi T.,et al.Mol. Cell. Biol. 12:836-846(1992).
Shen S.H.,et al.Nature 352:736-739(1991).
Shen S.H.,et al.Nature 353:868-868(1991).
Plutzky J.,et al.Proc. Natl. Acad. Sci. U.S.A. 89:1123-1127(1992).
Banville D.,et al.Genomics 27:165-173(1995).