

c-JUN Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52741**Specification**

c-JUN Antibody - Product Information

Application	WB
Primary Accession	P05412
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	43/48 KDa

c-JUN Antibody - Additional Information**Gene ID** 3725**Other Names**

Activator Protein 1;AP 1;AP1;cJun;Enhancer Binding Protein AP1;Jun Activation Domain Binding Protein;JUN;Jun oncogene;JUN protein;Jun proto oncogene;JUN_HUMAN;JUNC;Oncogene JUN;p39;Proto oncogene c jun;Proto oncogene cJun;Proto-oncogene c-jun;Transcription Factor AP 1;Transcription factor AP-1;Transcription Factor AP1;V jun avian sarcoma virus 17 oncogene homolog;V jun sarcoma virus 17 oncogene homolog (avian);V jun sarcoma virus 17 oncogene homolog;V-jun avian sarcoma virus 17 oncogene homolog;vJun Avian Sarcoma Virus 17 Oncogene Homolog.

Dilution

WB~~1:1000

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

c-JUN Antibody - Protein Information**Name** JUN**Function**

Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:10995748, PubMed:22083952). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:12618758). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:17210646). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:24623306). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:24623306).

Cellular Location

Nucleus.

Tissue Location


Expressed in the developing and adult prostate and prostate cancer cells.

c-JUN Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

c-JUN Antibody - Images



Western blot detection of c-JUN in CHO-K1 transfected by PEGFP-C1-c-JUN [A] and CHO-K1 cell lysate [B] cell lysate using c-JUN mouse mAb (1:1000 diluted). Predicted band size: 43/48KDa. Observed band size: 43/48KDa.

c-JUN Antibody - Background

Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'.

Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation.

c-JUN Antibody - References

Hattori K.,et al.Proc. Natl. Acad. Sci. U.S.A. 85:9148-9152(1988).
Bohmann D.,et al.Science 238:1386-1392(1987).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Gregory S.G.,et al.Nature 441:315-321(2006).