

Anti-Cytokeratin 18 Mouse mAb
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP53488**Specification**

Anti-Cytokeratin 18 Mouse mAb - Product Information

Application	WB, IHC
Primary Accession	P05783
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse IgG2b
Immunogen	Purified recombinant fragment of human Cytokeratin 18 (aa391-483) expressed in E. Coli.
Purification	Acites
Calculated MW	48kDa KDa
Antigen Region	aa391-483

Anti-Cytokeratin 18 Mouse mAb - Additional Information**Gene ID** 3875**Other Names**

CK18; CYK18; KRT18

Dilution

WB~~1:1000

IHC~~1:400

Format

Ascitic fluid containing 0.09% (W/V) sodium azide.

Storage

Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Anti-Cytokeratin 18 Mouse mAb - Protein Information**Name** KRT18**Synonyms** CYK18**Function**

Involved in the uptake of thrombin-antithrombin complexes by hepatic cells (By similarity). When phosphorylated, plays a role in filament reorganization. Involved in the delivery of mutated CFTR to the plasma membrane. Together with KRT8, is involved in interleukin-6 (IL-6)-mediated barrier protection.

Cellular Location

Nucleus matrix {ECO:0000250|UniProtKB:Q5BJY9}. Cytoplasm, perinuclear region. Nucleus, nucleolus. Cytoplasm {ECO:0000250|UniProtKB:Q5BJY9}


Tissue Location

Expressed in colon, placenta, liver and very weakly in exocervix. Increased expression observed in lymph nodes of breast carcinoma.

Anti-Cytokeratin 18 Mouse mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Cytokeratin 18 Mouse mAb - Images

Immunohistochemical analysis of KRT18 in Human appendix tissue sections(IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde at room temperature; antigen retrieval was by heat mediation with a EDTA buffer (pH9.0). Samples were incubated with primary antibody (1/400) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

Anti-Cytokeratin 18 Mouse mAb - Background

Cytokeratin 18, also known as CK18, CYK18, KRT18. Entrez Protein NP_000215. It encodes the type I intermediate filament chain keratin 18. Keratin 18, together with its filament partner keratin 8, are perhaps the most commonly found members of the intermediate filament gene family. They are expressed in single layer epithelial tissues of the body. Mutations in this gene have been linked to cryptogenic cirrhosis. Two transcript variants encoding the same protein have been found for this gene.